

RESEARCH ARTICLE

Evaluation of Peppermint (*Mentha piperita* L.) Essential Oil as a Digestive Tract Regulator in Broilers

Tuba BÜLBÜL¹  Vural ÖZDEMİR²  Aziz BÜLBÜL³ 

¹ Mugla Sıtkı Koçman University, Faculty of Milas Veterinary Medicine, Department of Animal Nutrition and Nutritional Disease, TR-48200 Muğla - TÜRKİYE

² Afyon Kocatepe University, Faculty of Veterinary Medicine, Department of Anatomy, TR-03200 Afyonkarahisar - TÜRKİYE

³ Mugla Sıtkı Koçman University, Faculty of Milas Veterinary Medicine, Department of Physiology, TR-48200 Muğla - TÜRKİYE



(*) **Corresponding author:** Tuba BÜLBÜL

Phone: +90 505 493 4572

E-mail: tubabulbul@mu.edu.tr

How to cite this article?

Bülbül T, Özdemir V, Bülbül A: Evaluation of peppermint (*Mentha piperita* L.) essential oil as a digestive tract regulator in broilers. *Kafkas Univ Vet Fak Derg*, 29 (5): 491-496, 2023.

DOI: 10.9775/kvfd.2023.29460

Article ID: KVFD-2023-29460

Received: 27.03.2023

Accepted: 11.09.2023

Published Online: 13.09.2023

ABSTRACT

The aim of this study was to investigate the effects of peppermint (*Mentha piperita* L.) essential oil on duodenal and colonic contractions of broilers *in vitro*. For this purpose, 10 broiler intestines (42 days old) obtained from a slaughterhouse were brought to the laboratories by immersed in Krebs solution (NaCl, 118 mmol/L; KCl, 4.7 mmol/L; CaCl₂, 2.5 mmol/L; MgSO₄, 1 mmol/L; KH₂PO₄, 1 mmol/L; glucose, 11 mmol/L; NaHCO₃, 25 mmol/L) at +4°C. Isolated duodenal and colon tissue strips were placed in a four-channel, isolated organ bath system, where they were exposed to Krebs solution aerated with a gas mixture of 95% O₂ - 5% CO₂, at 39°C. The effect of peppermint essential oil (PEO) obtained by hydrodistillation method on spontaneous duodenum and colon contractions was evaluated in the concentration range of 0.1-1000 µg/mL. Subsequently, the effects of 300 and 1000 µg/mL PEO were reevaluated on acetylcholine-induced contractions after incubation with N ω -Nitro-L-arginine (L-NNA, NOS inhibitor) and Methylene Blue (MB, cGC inhibitor). It was determined that the PEO dose-dependently decreased the amplitude of spontaneous and acetylcholine-induced contractions in both isolated tissues. This relaxant effect persisted after L-NNA and MB incubations. In conclusion, PEO induced relaxation in isolated duodenum and colon intestinal smooth muscles not through the nitric oxide-sGMP pathway.

Keywords: Broiler, Colon, Contraction, Duodenum, Peppermint essential oil

INTRODUCTION

Antibiotics are prohibited from being used as feed additives in poultry nutrition in the European Union because they cause a decrease in beneficial microorganisms in the intestinal microflora of animals, create residual risk in animal products, and adversely affect the health of people fed with these products [1]. Accordingly, studies on the development of alternative feed additives to antibiotics have been accelerated. In this context, the use of natural and reliable herbal products (aromatic plants and essential oils extracted from these plants) for poultry nutrition has gained importance [2,3]. Essential oils obtained by steam distillation or squeezing from the leaves, flowers, bark, seeds, and roots of plants have volatile and fragrant properties at room temperature [4]. Many studies have reported that these products protect the endocrine and immune systems [3,5], increase the digestibility and absorption of nutrients [6-8] and improve performance [9-11],

as well as antioxidant, anti-inflammatory, and antimicrobial effects [12].

Mentha plants are one of the most popular plants used since ancient times for their medicinal and aromatherapy properties [13]. The essential oil of peppermint, which is the most important *Mentha* specie, has been reported to have antioxidant, antidiabetic, antibacterial, anti-mutagenic, antifungal, and anticarcinogenic effects [14]. Studies conducted on the use of *Mentha* plants in poultry nutrition have evaluated the effects on performance [13,15], carcass traits [16,17], immunity, and some serum biochemistry parameters [18] during the growth period in broilers and on performance, egg quality and serum parameters in laying hens [19,20]. It is also stated that peppermint has a beneficial effect on the absorption surface area by improving the small intestine length and villi structure in quails [18]. These plants are used for the treatment of gastrointestinal (GI) system diseases [21-23].



The regulation of intestinal motility is significant as it adequate digestion of food and minimizes the risk of colonization by intestinal pathogens in the lower GI tract [24,25]. Prior studies have provided mixed results regarding the effect of essential oils and extracts derived from aromatic plants on intestinal contractions, as noticed in studies conducted on rats [22,23] and broilers [26]. This study was designed to evaluate the effect of various levels of POE on contractions occurring in the duodenum and colon of poultry. Assessing the potential impact of POE on contractions in the chicken GI tract will contribute to the development of potential uses for POE, which serves as an alternative to synthetic agents, such as antibiotics, in the field of poultry nutrition.

MATERIAL AND METHODS

Ethical Statement

This study was approved by the Afyon Kocatepe University Animal Experiments Local Ethics Committee (Approval no: AKÜ-HADYEK-272-13).

Extraction of PEO

Fresh mint (*Mentha piperita* L.) was obtained from a local herbal market in Afyonkarahisar, Turkey. After being cleared of weeds and parts, it was dried at room temperature without sunlight and then ground into powder using a mechanical grinder. 50 g of ground mint were transferred to a glass flask filled with 500 mL of distilled water. Hydrodistillation was carried out in a Clevenger apparatus for 180 min [26]. Before starting the distillation process, 1 mL of n-hexane was added to the water surface of the side arm of the Clevenger apparatus, and the condensed essential oil droplets were collected, reducing their distribution with water, and increasing efficiency. The resulting essential oil was dried over anhydrous Na₂SO₄ (Sigma-Aldrich, 238597) and stored in a closed dark glass bottle in a refrigerator at 4°C until use [25].

Animal and Tissue Preparation

This study was carried out on 10 broilers (Ross 308; 42 days old). They were obtained from the local private poultry slaughterhouse with standard regulations; the birds were sacrificed via a neck cut and bled for 120 sec. The birds were eviscerated manually [27]. The tissue strips were processed approximately 10 min after the animal was sacrificed, 2 cm-long strips of the mid-duodenum and colon were obtained. Then, these were immersed in a cold, freshly prepared Krebs solution (NaCl, 118 mmol/L; KCl, 4.7 mmol/L; CaCl₂, 2.5 mmol/L; MgSO₄, 1 mmol/L; KH₂PO₄, 1 mmol/L; glucose, 11 mmol/L; NaHCO₃, 25 mmol/L) and transferred to the laboratory. After removing fat and connective tissue from the intestinal segments,

longitudinal strips (0.6 cm long and approximately 3-4 mm wide) were placed in a four-chamber organ bath (IOBS 99 Isolated Tissue Bath Stand Set; Commat) with 20 mL Krebs solution (pH 7.4) in 95% O₂ and 5% CO₂ at 39°C. Longitudinal smooth muscle strips were attached to platinum ring electrodes along one edge using 2:0 silk ligatures. The opposite edge of the tissue was connected to a force-displacement transducer (model 10-A; MAY; Commat, Ankara, Türkiye). The isometric smooth muscle activity of the intestine samples was monitored and recorded by the computer using the force transducer and an acquisition system (model MP30 WSW with Biopac Student Lab, PRO Software, Biopac Systems; Commat). To maintain the contractile activity of the tissue, optimal tension relationships for the strips were achieved with resting tensions of 1 g. It was allowed to equilibrate for an hour and the Krebs solution was changed every fifteen minutes. The viability of the strips was verified by adding Acetylcholine (Ach) at a concentration of 10⁻⁵ M at the beginning and/or end of the experiment.

Monitoring the GI Tract Muscle Activity

Subsequently, the effectiveness of the doses (300 and 1000 µg/mL) of PEO that cause inhibition in duodenal and colonic contractions was reevaluated after incubation with the NOS inhibitor L-NNA (10⁻⁵ M) and the cGC blocker MB (10⁻² M). In addition, the effect of 8-bromocyclic GMP (8-Br-cGMP; 10⁻⁸, 10⁻⁷ and 10⁻⁶ M), a cGMP analog, on the amplitude of isolated duodenal and colon spontaneous contractions was evaluated.

Statistical Analysis



Statistical analysis was done using SPSS 22.0 (SPSS Inc., Chicago, IL, USA). The Shapiro-Wilk test was used to test the data for normality. All values are presented as mean ± S.E.M. One-way ANOVA was used for the statistical evaluation of the data. The Tukey test was performed to compare individual means of treatment groups. Differences were considered significant at P<0.05.

RESULTS

Effect of PEO on the Amplitude of Spontaneous Duodenum and Colon Contractions

It was determined that the cumulative application of PEO at concentrations of 0.1, 0.3, 1, 3, 10, 30 100, 300 and 1000 µg/mL caused inappropriate relaxation to the gradual concentration response in the duodenum (*Fig. 1-A*) and colon (*Fig. 1-B*) tissue samples.

The amplitude of spontaneous duodenal and colonic contractions was reduced by PEO at concentrations of 300 µg/mL (P<0.01) and 1000 µg/mL (P<0.001). However, no statistically significant influence on these contractions was seen at concentrations from 0.1 to 100 µg/mL. It was



observed that the contractions returned to normal after washing the tissues with Krebs solution.

Effect of PEO on Ach-Induced Duodenum and Colon Contractions

The effects of 300 and 1000 µg/mL PEO on duodenal and colonic contractions induced by Ach (10^{-4} M), respectively, are shown in Fig. 2. It was determined that 1000 µg/mL PEO reduced the contraction effect of Ach in both tissues ($P < 0.05$).

Effect of PEO on Duodenum and Colon Contractions Induced by Ach Following L-NNA and MB Incubation

The effects of 300 and 1000 µg/mL PEO on the amplitude

of duodenal and colonic contractions induced by Ach (10^{-4} M) following L-NNA and MB incubations, respectively, are given in Table 1 and Table 2. It was determined that 1000 µg/mL PEO decreased the amplitude of contraction in both tissues ($P < 0.05$), and a similar effect was observed after MB incubation ($P < 0.05$).

Effect of 8-Br-cGMP on Spontaneous Duodenum and Colon Contractions

The effect of 8-Br-cGMP (10^{-8} - 10^{-6} M) on spontaneous duodenal and colonic contractions is shown in Fig. 3. It was observed that 10^{-8} M dose of 8-Br-cGMP had no effect on the contraction intensity of both tissues, whereas 10^{-7}

Tissue	L-NNA + Ach	L-NNA + Ach + PEO 300	L-NNA ± Ach + PEO 1000	P-value
Duodenum	0.661±0.052 ^a	0.608±0.036 ^{ab}	0.511±0.026 ^b	0.048
Colon	1.301±0.151 ^a	1.190±0.135 ^{ab}	0.955±0.092 ^b	0.032

L-NNA: Nω-Nitro-L-arginine; Ach: Acetylcholine; PEO: peppermint essential oil
Letters (a, b) in the same line indicate significant differences between different letters

Tissue	MB + Ach	MB + Ach + PEO 300	MB ± Ach + PEO 1000	P-value
Duodenum	0.676±0.053 ^a	0.608±0.037 ^a	0.496±0.021 ^b	0.010
Colon	1.313±0.152 ^a	1.208±0.146 ^{ab}	0.945±0.093 ^b	0.042

MB: Methylene Blue; Ach: Acetylcholine; PEO: peppermint essential oil
Letters (a, b) in the same line indicate significant differences between different letters




Fig 3. The dose-response curves of 8-Br-cGMP in isolated broiler duodenum and colon (A: Duodenum, B: Colon). Letters (a, b) in the same line indicate significant differences between different letters

M and 10⁻⁶ M doses depressed the contraction intensity ($P < 0.001$).

DISCUSSION

In this study, PEO decreased the amplitude of spontaneous duodenum and colon contractions in broilers in a dose-dependent manner. Similarly, we have previously demonstrated that myrtle, rosemary, and thyme essential oils inhibit contractions of the small intestine [26]. Essential oils have been added to poultry diets for several purposes. Recently, they have also been used to increase the digestibility and absorption of nutrients in poultry [7,28,29]. Essential oils have been demonstrated to increase the secretion of some digestive enzymes [6] and stimulate the appetite center due to their aroma-enhancing properties [30,31]. Since essential oils obtained from different plants regulate the contractions in the GI tract of rodents, the use of these plants is recommended to support the digestive tract [32,33]. Mentha plants have been used in traditional medicine for many years in the treatment of GI disorders [13]. In this study, the spasmolytic effect created by the essential oil of peppermint, known as *Mentha spicata*, was demonstrated in isolated duodenum and colon smooth muscle.

In the GI tract, Ach is an important neural mediator of parasympathetic innervation in poultry, similar to in other animals. It binds to muscarinic receptors and induces smooth muscle contraction [34,35]. Our previous study showed which Ach concentrations of 10⁻⁴ M induce maximal contractions in the duodenum of broilers [26]. In this study, duodenum and colon contractions were induced by using Ach (10⁻⁴ M), and the effectiveness of PEO (300-1000 g/mL) was reevaluated. The spasmolytic activity of PEO has been demonstrated by reducing the amplitude of Ach-induced contractions, similar to spontaneous duodenal and colonic contractions.

In this study, after determining the relaxation effect of PEO, we hypothesized that it may be efficient through the NOS-NO-cGC. The smooth muscle layers are innervated by enteric inhibitory and excitatory motor neurons that directly regulate GI motility [36]. Nitric oxide (NO), which is released in response to nerve stimulation of the

myenteric plexus, causes relaxation of smooth muscle in addition to the protection of the digestive tract mucosa as a result of the regulation of gastric mucosal blood flow and fluid secretion. Thus, it plays an important role in regulating stomach and intestinal motility [37,38]. In the myenteric plexus, neuronal NO synthase (nNOS) generates NO from L-arginine [36]. Due to their ability to inhibit NOS, L-arginine analogs, specifically N-nitro-L-arginine (L-NNA) and N- ω -Nitro-L-arginine methyl ester, are frequently used to prevent the production of NO [40,41]. In contraction trials, when the nNOS inhibitor L-NNA is at a concentration of 1 mM, it blocks the NOS enzyme [41,42]. In the present study, it was determined that the relaxing effect of PEO continued in the duodenum and colon, induced by Ach, after inhibition of the NOS enzyme with L-NNA. Therefore, the results of the study show that PEO does not create a relaxant effect by increasing NO formation.

The primary receptor for NO is soluble guanylate cyclase (sGC) [40]. Nitric oxide activation of sGC leads to the formation of cGMP. As a second messenger, cGMP interacts with protein kinases, phosphodiesterases, and ion channels, leading to the cellular effects of NO. The cGMP induces relaxation in smooth muscle tissue by two mechanisms [44]. First, it increases the permeability of K⁺ channels by lowering the level of Ca⁺⁺ in the cell. Thus, it hyperpolarizes the plasma membrane [45,46]. Second, cGMP blocks the myosin/actin interaction by activating cGMP-dependent protein kinase (PKG), which causes the dephosphorylation of myosin light chains [47]. Moreover, some of the NO-induced smooth muscle relaxations were also reported to be independent of cGMP [36,48]. However, there are no previous studies in which the effects and dose were demonstrated in the anatomical formations of the small and large intestines, although the effect of cGMP was shown in different smooth muscle cells in poultry. In this study, it was determined that the amplitude of duodenum and colon contractions decreased. In this context, a cGMP-dependent relaxation was observed in the poultry intestine.

Methylene blue prevents cGMP-dependent relaxation by inhibiting cGC inhibitor and cGMP formation [49]. It is known that the dose of 10⁻⁵ M blocks cGC in *in vitro*

experiments ^[42,50]. The present study provided evidence that the intensity of Ach-induced duodenal and colonic contractions was reduced by PEO after inhibition of MB and cGC, whereas MB did not reduce PEO activity. In conclusion, it is considered that the effectiveness of PEO is not a cGMP dependent.

As a result, it was determined that a cGMP-dependent relaxation was observed in the duodenum and colon of broilers, whereas PEO reduced the amplitude of contractions in the same tissues without using the NOS-cGC-cGMP pathway. It is suggested that >10 µg/mL of PEO be added to broiler diets due to its spasmolytic effect, but it is also recommended to conduct trials in terms of *in vivo* activity.

Availability of Data and Materials

The datasets used and/or analyzed during the current study are available from the corresponding author (T. Bülbül) on reasonable request.

Funding Support

This study was supported by the Scientific Research Projects Unit at Afyon Kocatepe University under project number 17.KARIYER.183.

Ethical Statement

This study was approved by the Afyon Kocatepe University Animal Experiments Local Ethics Committee (Approval no: AKÜ-HADYEK-272-13).

Conflict of Interest

There is no conflict of interest.

Author Contributions

TB designed this study, prepared this manuscript, and analyzed experimental data; VÖ analyzed experimental data; AB analyzed experimental data.

REFERENCES

- Jankowski J, Zdunczyk Z, Juskiewicz J, Kozłowski K, Leczewicz A, Jeroch H: Gastrointestinal tract and metabolic response of broilers to diets with the *Macleaya cordata* alkaloid extract. *Arch Fur Geflügelkunde*, 73, 95-101, 2009.
- Zhai H, Liu H, Wang S, Wu J, Klünter AM: Potential of essential oils for poultry and pigs. *Anim Nutr*, 4 (2): 179-186. 2018. DOI: 10.1016/j.aninu.2018.01.005
- Puvača N, Tufarelli V, Giannenas I: Essential oils in broiler chicken production, immunity and meat quality: Review of *Thymus vulgaris*, *Origanum vulgare*, and *Rosmarinus officinalis*. *Agriculture*, 12 (6): 874, 2022. DOI: 10.3390/agriculture12060874
- Dhifi W, Bellili S, Jazi S, Bahloul N, Mnif W: Essential oils chemical characterization and investigation of some biological activities: A critical review. *Medicines*, 3 (4):25, 2016. DOI: 10.3390/medicines3040025
- Francois R: Active plant extracts show promise in poultry production. *Poult Int*, 20, 28-31, 2006.
- Lee KW, Everts H, Kappert HJ, Frehner M, Losa R, Beynen AC: Effects of dietary essential oil components on growth performance, digestive enzymes and lipid metabolism in female chickens. *Br Poult Sci*, 44 (3): 450-457, 2003. DOI: 10.1080/0007166031000085508
- Platel K, Srinivasan K: Digestive stimulant action of spices: A myth or reality. *Indian J Med Res*, 119, 167-179, 2004.
- Jamroz D, Wertelecki T, Houszka M, Kamel C: Influence of diet type on the inclusion of plant origin active substances on morphological and histochemical characteristics of the stomach and jejunum walls in chicken. *J Anim Physiol Anim Nutr*, 90, 255-268, 2006. DOI: 10.1111/j.1439-0396.2005.00603.x
- Ölmez M, Adıgüzel Işık S, Şahin T, Yörük MA: The effects of pennyroyal (*Mentha pulegium* L.) on performance, carcass and meat quality in Japanese quails (*Coturnix coturnix japonica*). *Ankara Univ Vet Fak Derg*, 68 (4): 343-348, 2021. DOI: 10.33988/aufvd.767472
- Biricik H, Yesilbag D, Gezen SS, Bulbul T: Effects of dietary myrtle oil (*Myrtus communis* L.) supplementation on growth performance, meat oxidative stability, meat quality and erythrocyte parameters in quails. *Revue Med Vet*, 3, 131-138, 2012.
- Yesilbag D, Gezen SS, Biricik H, Bulbul T: Effect of a rosemary and oregano volatile oil mixture on performance, lipid oxidation of meat and haematological parameters in pharaoh quails. *Br Poult Sci*, 53, 89-97, 2012. DOI: 10.1080/00071668.2012.654763
- Gharaibeh MH, Khalifeh MS, Nawasreh AN, Hananeh WM, Awawdeh MS: Assessment of immune response and efficacy of essential oils application on controlling necrotic enteritis induced by clostridium perfringens in broiler chickens. *Molecules*, 26 (15): 4527. 2021. DOI: 10.3390/molecules26154527
- Sameer Shaikh S: Prospective role in treatment of major illnesses and potential benefits as a safe insecticide and natural food preservative of mint (*Mentha* spp.): A review. *Asian J Biomed Pharm Sci*, 4, 1-12, 2014. DOI: 10.15272/ajbps.v4i35.559
- Amani Machiani M, Javanmard A, Morshedloo MR, Maggi F: Evaluation of yield, essential oil content and compositions of peppermint (*Mentha piperita* L.) intercropped with faba bean (*Vicia faba* L.). *J Clean Prod*, 171, 529-537, 2018. DOI: 10.1016/j.jclepro.2017.10.062
- Gürbüz Y, İsmail IA: Effect of peppermint and basil as feed additive on broiler performance and carcass characteristics. *Iran J Appl Anim Sci*, 6 (1): 149-156, 2015.
- Khempaka S, Pudpila U, Molee W: Effect of dried peppermint (*Mentha cordifolia*) on growth performance, nutrient digestibility, carcass traits, antioxidant properties, and ammonia production in broilers. *J Appl Poult Res*, 22, 904-912, 2013. DOI: 10.3382/japr.2013-00813
- Mehri M, Sabaghi V, Bagherzadeh-Kasmani F: *Mentha piperita* (peppermint) in growing Japanese quails diet performance, carcass attributes, morphology and microbial populations of intestine. *Anim Feed Sci Technol*, 207, 104-111, 2015. DOI: 10.1016/j.anifeeds.2015.05.021
- Mehri M, Sabaghi V, Bagherzadeh-Kasmani F: *Mentha piperita* (peppermint) in growing Japanese quails' diet serum biochemistry, meat quality, humoral immunity. *Anim Feed Sci Technol*, 206, 57-66, 2015. DOI: 10.1016/j.anifeeds.2015.05.022
- Paymard J, Nobakht A, Mazlum F, Moghaddam M: The effects of different levels of dried aerial parts powder and extract of pennyroyal (*Mentha pulegium*) medicinal plant on performance, egg quality, blood biochemical and immunity parameters of laying hens. *Iran J Appl Anim Sci*, 3 (3): 589-594, 2013.
- Abdel-Wareth AAA, Lohakare JD: Effect of dietary supplementation of peppermint on performance, egg quality, and serum metabolic profile of hy-line brown hens during the late laying period. *Anim Feed Sci Technol*, 197, 114-120, 2014. DOI: 10.1016/j.anifeeds.2014.07.007
- Tayarani-Najaran Z, Talasaz-Firooz E, Nasiri R, Jalali N, Hassanzadeh MK: Antiemetic activity of volatile oil from *Mentha spicata* and *Mentha × piperita* in chemotherapy induced nausea and vomiting. *Ecancermedicalscience*, 7:290, 2013. DOI: 10.3332/ecancer.2013.290
- Estrada-Soto S, Gonzalez-Maldonado D, Castillo-Espana P, Aguirre-Crespo F, Sanchez Salgado JC: Spasmolytic effect of *Mentha pulegium* L. involves ionic flux regulation in rat ileum trips. *J Smooth Muscle Res*, 46 (2): 107-117, 2010. DOI: 10.1540/jsmr.46.107
- Naseri MKG, Naseri ZG, Mohammadian M, Birgani MO: Ileal relaxation induced by *Mentha longifolia* (L.) leaf extract in rat. *Pak J Biol Sci*, 11 (12): 1594-1599, 2008. DOI: 10.3923/pjbs.2008.1594.1599
- Waclawiková B, Codutti A, Alim K, El Aidi S: Gut microbiota-motility interregulation: insights from *in vivo*, *ex vivo* and *in silico* studies. *Gut*

Microbes, 14 (1):1997296, 2022. DOI: 10.1080/19490976.2021.1997296

25. **Tang L, Cheng CY, Sun X, Pedicone AJ, Mohamadzadeh M, Cheng SX:** The extracellular calcium-sensing receptor in the intestine: evidence for regulation of colonic absorption, secretion, motility, and immunity. *Front Physiol*, 21 (7):245, 2016. DOI: 10.3389/fphys.2016.00245

26. **Bülbül T, Özdemir V, Ulutaş E, Bülbül A:** Broiler tavuklarda mersin, biberiye ve kekik esansiyel yağlarının bağırsak motilitesi üzerine etkisi. *Kocatepe Vet J*, 11 (4): 394-401, 2018. DOI: 10.30607/kvj.434247

27. **Aksu T, Aksu MI, Yoruk MA, Karaoglu M:** Effects of organically-complexed minerals on meat quality in chickens. *Br Poult Sci*, 52 (5): 558-63, 2011. DOI: 10.1080/00071668.2011.606800

28. **Burtis M, Bucar F:** Antioxidant activity of *Nigella sativa* essential oil. *Phytother Res*, 14, 323-328, 2000. DOI: 10.1002/1099-1573(200008)14:5<323::aid-ptr621>3.0.co;2-q

29. **Hernandez F, Madrid J, Garcia V, Orengo J, Megias MD:** Influence of two plant extracts on broilers performance, digestibility and digestive organ size. *Poult Sci*, 83, 169- 174, 2004. DOI: 10.1093/ps/83.2.169

30. **Deyoe CW, Davies RE, Krishnan R, Khaund R, Couch JR:** Studies on the taste preference of the chick. *Poult Sci*, 41, 781-784, 1962. DOI: 10.3382/ps.0410781

31. **Te Pas MFW, Borg R, Buddiger NJH, Wood BJ, Rebel JMJ, van Krimpen MM, Calus MPL, Park JE, Schokker D:** Regulating appetite in broilers for improving body and muscle development - A review. *J Anim Physiol Anim Nutr (Berl)*, 4 (6): 1819-1834, 2020. DOI: 10.1111/jpn.13407

32. **Sadraei H, Ghannadi A, Malekshahi K:** Relaxant effect of essential oil of *Melissa officinalis* and citral on rat ileum contractions. *Fitoterapia*, 74, 445-452, 2003. DOI: 10.1016/s0367-326x(03)00109-6

33. **Ghader JA, Massoud M, Bahram DN, Farshad K:** Effects of *Bunium persicum* (boiss.) essential oil on the contractile responses of smooth muscle: An *in vitro* study. *Vet Res Forum*, 2, 87-96, 2011.

34. **Tanahashi Y, Komori S, Matsuyama H, Kitazaw T, Unno T:** Functions of muscarinic receptor subtypes in gastrointestinal smooth muscle: A review of studies with receptor-knockout mice. *Int J Mol Sci*, 2, 926, 2021. DOI: 10.3390/ijms22020926

35. **Iino S, Ward SM, Sanders KM:** Interstitial cells of Cajal are functionally innervated by excitatory motor neurones in the murine intestine. *J Physiol*, 556, 521-530, 2004. DOI: 10.1113/jphysiol.2003.058792

36. **Idrizaj E, Traini C, Vannucchi MG, Baccari MC:** Nitric oxide: From gastric motility to gastric dysmotility. *Int J Mol Sci*, 22, 9990, 2021. DOI: 10.3390/ijms22189990

37. **Goyal RK, Guo Y, Mashimo H:** Advances in the physiology of gastric emptying. *Neurogastroenterol Motil*, 31:e13546, 2019. DOI: 10.1111/nmo.13546

38. **Wallace JL:** Nitric oxide in the gastrointestinal tract: opportunities for drug development. *Br J Pharmacol*, 176 (2): 147-154, 2019. DOI: 10.1111/bph.14527

39. **Vannucchi, MG, Corsani L, Bani D, Fausone-Pellegrini MS:** Myenteric neurons and interstitial cells of Cajal of mouse colon express several nitric oxide synthase isoforms. *Neurosci Lett*, 326, 191-195, 2002. DOI: 10.1016/s0304-3940(02)00338-5

40. **Groneberg D, Voussen B, Friebe A:** Integrative control of gastrointestinal motility by nitric oxide. *Curr Med Chem*, 23, 2715-2735, 2016. DOI: 10.2174/0929867323666160812150907

41. **Ren H, Yuan F, Tan W, Ding Y, An P, Luo H:** Effect of evodiamine on rat colonic hypermotility induced by water avoidance stress and the underlying mechanism. *Drug Des Devel Ther*, 15, 441-452, 2021. DOI: 10.2147/DDDT.S298954

42. **Bulbul A, Yagci A, Altunbas K, Sevimli A, Celik HA, Karadeniz A, Akdag E:** The role of nitric oxide in the effects of ovarian steroids on spontaneous myometrial contractility in rats. *Theriogenology*. 68 (8): 1156-1168, 2007. DOI: 10.1016/j.theriogenology.2007.08.014

43. **Friebe A, Voußen B, Groneberg D:** NO-GC in cells 'off the beaten track. *Nitric Oxide*, 77, 12-18, 2018. DOI: 10.1016/j.niox.2018.03.020

44. **Mujoo K, Krumenacker JS, Murad F:** Nitric oxide-cyclic GMP signaling in stem cell differentiation. *Free Radic Biol Med*, 51 (12): 2150-2157, 2011. DOI: 10.1016/j.freeradbiomed.2011.09.037

45. **Jackson WF:** Endothelial ion channels and cell-cell communication in the microcirculation. *Front Physiol*, 13:805149, 2022. DOI: 10.3389/fphys.2022.805149

46. **Valero MS, López V, Castro M, Gómez-Rincón C, Arruebo MP, Les E, Plaza MÁ:** Involvement of NO/cGMP signaling pathway, Ca²⁺ and K⁺ channels on spasmolytic effect of everlasting flower polyphenolic extract (*Helichrysum stoechas* (L.) Moench). *Int J Mol Sci*, 23 (22): 14422, 2022. DOI: 10.3390/ijms232214422

47. **Francis SH, Busch JL, Corbin JD, Sibley D:** cGMP-dependent protein kinases and cGMP phosphodiesterases in nitric oxide and cGMP action. *Pharmacol Rev*, 62 (3): 525-63, 2010. DOI: 10.1124/pr.110.002907

48. **Martins SLR, De Oliveira RB, Ballejo G:** Rat duodenum nitregic-induced relaxations are cGMP-independent and apamin-sensitive. *Eur J Pharmacol*, 284, 265-270, 1995. DOI: 10.1016/0014-2999(95)00348-o

49. **Ginimuge PR, Jyothi SD:** Methylene blue: Revisited. *J Anaesthesiol Clin Pharmacol*, 26 (4): 517-520, 2010.

50. **Jing M, Ling GS, Bina S, Hart JL, Muldoon SM:** Halothane attenuates nitric oxide relaxation of rat aortas by competition for the nitric oxide receptor site on soluble guanylyl cyclase. *Eur J Pharmacol*, 342 (2-3): 217-224, 1998. DOI: 10.1016/s0014-2999(97)01491-x