



Research Article

LIPID COMPOSITIONS OF *Bjerkandera adusta* (WILLD) P KARST

Selçuk KÜÇÜKAYDIN¹, Mehmet Emin DURU^{2*}

¹Department of Chemistry, Faculty of Science, Mugla Sıtkı Kocman University, MUĞLA;
ORCID:0000-0001-8538-6528

²Department of Chemistry, Faculty of Science, Mugla Sıtkı Kocman University, MUĞLA;
ORCID:0000-0001-7252-4880

Received: 25.02.2017 Revised: 17.03.2017 Accepted: 10.04.2017

ABSTRACT

Mushrooms are considered as a substantial nutrient, which are the source of many vitamins as well as protein and mineral. Since ancient times, it has been known that mushrooms used as food as well as medicine in many societies. In addition to the nutritional properties of mushrooms, it is especially preferred by people due to its flavor and aroma. Despite the low fat content of mushrooms, the high proportion of polyunsaturated fatty acids further increasing the importance of mushrooms for nutrition. In this study, fatty acid components of an edible mushroom *Bjerkandera adusta* was investigated. It was determined that 20 of the 26 components were fatty acid in the lipid fraction of *B. adusta* by using GC and GC/MS systems. Linoleic acid (37.25%), oleic acid (9.51%) and stearic acid (4.51%) were found as major fatty acids of *B. adusta*. Additionally, ergosterol (12.46%) was found in high concentration in the lipid fraction of *B. adusta*. As a result, it was found that the percentage of unsaturated fatty acids in the total lipid fraction is 52.69% in *B. adusta*.

Keywords: *Bjerkandera adusta*, fatty acid, ergosterol, GC, GC/MS, mushroom.

1. INTRODUCTION

In the past, mushrooms have known as a “plant”, but later on it has recognized separate kingdom due to its different characters from plants and animals. Although, many mushroom species are poisonous, some of them have medicinal usage. Some of the mushroom species particularly known as “parasol mushrooms” are used as food. [1-3]. Mushrooms are widely consumed as a food in many countries and it is one of the essential foods due to its impressive tastes, aromas and nutritional properties. Mushrooms are valuable food items due to having high protein, fiber, vitamins, minerals and low fat content [4-6].

Mushrooms have attracted the attention of researchers in recent years, according to their biological activities. Such as; The immunomodulatory, antitumor as well as antiviral, antimicrobial, antimutagenic, antihypertensive, antiinflammatory, antiallergic activities are besides the biological activities of mushrooms [7-8].

Up to date mainly, Lecthin, polysaccharides, polysaccharide-peptides, polysaccharide-protein types of compounds have been isolated from the mushrooms which are these substances which are exhibiting immunomodulatory, anticancer and antioxidant activities [7-9].

* Corresponding Author/Sorumlu Yazar: e-mail/e-ileti: eminduru@mu.edu.tr, tel: (252) 211 14 94

In addition to medicinal properties of mushrooms, there is an increasing of importance on the utilization of them as a “functional food” due to their unique composition including proteins, amino acids, vitamins, minerals and high-content polyunsaturated fatty acid. Although the fat content of mushrooms account about 5%, and 75% of it consists of unsaturated fatty acid components, that makes mushrooms significant part of healthy nutrition. It is known that, especially polyunsaturated fatty acids, omega-3, and -6 series are necessary for human health in the prevention and treatment of hypertension, coronary artery disease, diabetes, osteoporosis, arthritis, cancer, other inflammatory and autoimmune disorders. Due to the medical properties of polyunsaturated fatty acids and their role in natural nutrition, scientific studies on mushrooms are increasing day by day [10-17].

In the literature, there is a short report on fatty acid composition of some mushroom species which are growing on permafrost conditions. According to that paper, linoleic acid, eicosanoic acid, oleic acid and stearic acid have been found major components of *B. adusta* respectively [18].

The aim of this work was to investigate the lipid composition of *Bjerkandera adusta* which is edible and growing naturally in Turkey using by GC-FID and GC-MS systems.

2. MATERIALS AND METHODS

2.1. Mushroom Material

B. adusta is growing naturally in Turkey and known as tree mushrooms due to, growing on tree barks. *B. adusta* was collected from sweetgum (*Liquidambar orientalis*) forests in Köyceğiz-Muğla, province of Turkey in 2015. They were identified and deposited in the Fungarium of Department of Biology, Muğla Sıtkı Koçman University and were stored at -18 °C until analyses (Fungarium no: AT-2447).

2.2. Chemicals and Instruments

Boron trifluoride-methanol solution (BF₃:MeOH), n-hexane and chloroform were obtained from E. Merck (Darmstadt, Germany). GC-FID analyses were performed on a Shimadzu GC-17 AAF, V3, 230 V series gas chromatography (Japan) and GC-MS analyses were on Varian Saturn 2100 (USA).

2.3. Extraction

Dry mushroom sample (50 g) was extracted with 100 mL n-hexane:chloroform (90:10) solvent system for three times (24 h x 3) at room temperature, then filtered and evaporated to dryness under vacuum. They stored at refrigerator (+4°C) until methylation process.

2.4. Methylation of lipid extract of *B. adusta*

The lipid extract (100 mg) was dissolved in 0.5 M NaOH (2 mL). After that solution was heated in a water bath (50 °C), then 2 mL BF₃:MeOH was added. The mixture was boiled for 3 minutes, and then the mixture was left until it cooled down, and then the volume was completed to 25 mL with saturated NaCl solution. Esters were extracted with n-hexane; thus, the organic layer was separated. The hexane layer was dried with anhydrous Na₂SO₄ and filtered. The organic solvent was removed under reduced pressure by a rotary evaporator to give methyl esters.

2.5. Gas chromatography (GC) Analysis

GC analysis of the methyl derivatives of lipid fraction of *B.adusta* was performed by Shimadzu GC-17 AAF, V3, 230 V series gas chromatography (Japan) coupled with a Flame Ionisation Detector (FID) and a DB-1 fused silica capillary non-polar column (30 m x 0.25 id., film thickness 0.25 µm). Injector and detector temperatures were 250 and 270 °C, respectively, carrier gas was He at a flow rate of 1.4 mL/min; sample volume, 0.4 µL; split ratio, 50:1. The initial oven temperature was held at 100 °C for 5 min, then increased up to 238 °C with 3 °C/min increments and held at this temperature for 9 minutes. The relative percentages of separated compounds were calculated by using GC Solution computer program.

2.6. Gas chromatography-Mass spectrometry (GC-MS) Analysis

GC-MS analysis of the methyl derivatives of lipid fraction determined using by Varian Saturn 2100T (USA) coupled with an ion trap mass spectrometer and a DB-1 MS fused silica non-polar capillary column (30 m x 0.25 mm ID, film thickness 0.25 µm). For GC-MSD detection, an electron ionization system with ionization energy of 70 eV was used. Carrier gas was helium (15 psi) at a flow rate of 1.3 mL/min. Injector and MS transfer line temperatures were set at 250 and 200 °C, respectively. The oven temperature was held at 100 °C for 5 min, then increased up to 238 °C with 3 °C/min increments and held at this temperature for 9 minutes. 0.2 µL was injected manually in the split mode. Split ratio was 50:1. EI-MS were taken at 70 eV ionization energy. Mass range was from m/z 28 to 650 amu. The library search was carried out using NIST and Wiley 2005 (Gas Chromatography-Mass Spectrometry) GC-MS libraries. FAME (Fatty acid Methyl Ester) mixture (Supelco™ 37, Catalog no: 47885-U) were identified by comparing their retention times with those of the pure FAMES standards.

3. RESULTS AND DISCUSSIONS

The yield of lipid fraction was found 2.50% in *B. Adusta*. The presence of twenty six components were determined in the lipid compositions of *B.adusta* (Table 1). Linoleic acid (C18:2 ω6) (37.25%), ergosterol (12.46%) and oleic acid (C18:1 ω9) (9.51%) were the major lipid components of *B.adusta*. The lipid components in the mushroom were observed to be rich in fatty acids and unsaturated fatty acid amounts were higher than saturated ones in *B.adusta*. Linoleic acid and oleic acid are very important for human diet. These compounds are known to be decreasing the risk of cardiovascular disease and also recommended for the balancing high blood cholesterol.

Besides fatty acids, Ergosterol (12.46%) was one of the major components. Ergosterol is known as provitamin D₂ and mostly found in mushrooms because it is a part of their cytoplasmic membrane. Ergosterol is converted to vitamin D₂ by exposure to ultraviolet (UV) light. Vitamin D₂ is a type of vitamin D and used as prevent and treat vitamin D deficiency. Vitamin D is important for dietary supplement as a regulator of the metabolism of calcium and phosphate [19]. There is some reports on medicinal activities of ergosterol such as; growth-inhibition of bladder cancer, lipid peroxidation, antihyperlipidemic and anti-inflammatory activities [20-23].

We found also unusual components for mushrooms such as, cinnamic acid (5.13%) and cinnamyl alcohol (0.65%). On the other hand, cinnamic acid derivatives are major constituents of resin of *Liquidambar orientalis* [24]. The *L. orientalis* (oriental sweetgum) is a tertiary period relict endemic taxon of the east Mediterranean [25]. Due to, *B.adusta* is growing on *Liquidambar orientalis*, cinnamic acid and cinnamyl alcohol might be derived from *Liquidambar orientalis*.

Table 1. Lipid Composition (%) of *Bjerkandera adusta*

Peak no	Compound	Concentration (%)
1	Cinnamyl alcohol	0.65
2	Mandelic acid	0.57
3	Cinnamic acid	5.13
4	9-oxo-nonanoic acid (C _{9,0})	0.63
5	10-methyl-undecanoic acid (C _{11,0})	0.34
6	2-oxo-cyclohexane propanoic acid	1.32
7	12-Methyl, tridecanoic acid (C _{13,0})	1.51
8	10-Pentadecen-1-ol	0.26
9	Pentadecanoic acid (C _{15,0})	3.60
10	Palmitic acid (C _{16,0})	3.14
11	Methyl, 3- (3,5-ditertbutyl-4-hydroxy-phenyl) propionate	1.23
12	3,15-Octadecadien-1-ol acetate	1.18
13	Margaric acid (C _{17,0})	0.83
14	Linoleic acid (C_{18,2})	37.25
15	Oleic acid (C_{18,1})	9.51
16	Stearic acid (C _{18,0})	4.51
17	Eicosanoic acid (C _{20,0})	0.88
19	Heneicosanoic acid (C _{21,0})	0.37
18	Docosatetraenoic acid (C _{22,4})	2.79
20	Behenic acid (C _{22,0})	1.65
21	Diisooctyl phthalate	1.35
22	Tricosanoic acid (C _{23,0})	1.31
23	15-tetracosanoic acid (C _{24,1})	3.14
24	Tetracosanoic acid (C _{24,0})	2.89
25	Ergosterol	12.46
26	Pentacosanoic acid (C _{25,0})	1.50
	Saturated fatty acids	23.16
	Unsaturated fatty acids	52.69
	Other components	24.15

4. CONCLUSIONS

Mushrooms are the valuable and healthy foods in human diet, due to their low-fat composition, low calories and high essential fatty acid levels. The present study indicates that *B.adusta* contains high percentage of unsaturated fatty acids especially linoleic acid and oleic acid. Mushrooms are known to be a good source of vitamin D₂. It contains high percentage of ergosterol (provitamin D₂) that has medicinal properties and also phenolic organic compounds were found such as cinnamic acid and cinnamyl alcohol in the lipid fraction of *B.adusta* as well. These results showed that, *B.adusta* has a great potential to use as a supplementary food or nutraceuticals.

Acknowledgements

This study is a part of Ph.D. thesis of S.Kucukaydin (Graduate School of Natural and Applied Sciences, Muğla Sıtkı Koçman University-Turkey).

REFERENCES

- [1] Kavishree, S., Hemavathy, J., Lokesh, B. R., Shashirekha, M. N., & Rajarathnam, S. (2008). Fat and fatty acids of Indian edible mushrooms. *Food Chemistry*, 106, 597.
- [2] Reis, F. S., Barros, L., Martins, A., & Ferreira, I. C. F. R. (2012). Chemical composition and nutritional value of the most widely appreciated cultivated mushrooms, an inter-species comparative study. *Food and Chemical Toxicology*, 50, 191.
- [3] Ozturk, M., Tel, G., Ozturk, F. A., & Duru, M. E. (2014). The Cooking Effect on Two Edible Mushrooms in Anatolia, Fatty Acid Composition, Total Bioactive Compounds, Antioxidant and Anticholinesterase Activities. *Records of Natural Products*, 8, 189.
- [4] De Pinho, P.G., Ribeiro, B., Gonçalves, R.F., Baptista, P., Valentao, P., Seabra, R.M. ve Andrade, P.B. (2008) Correlation between the pattern volatiles and the overall aroma of wild edible mushrooms, *J Agric Food Chem*, 56:1704-1712.
- [5] Barros, L., Cruz, T., Baptista, P., Estevinho, L.M. ve Ferreira, I.C.F.R. (2008) Wild and commercial mushrooms as source of nutrients and nutraceuticals, *Food Chem Toxicol*, 46: 2742-2747.
- [6] Ribeiro, B., de Pinho, P. G., Andrade, P. B., Baptista, P., & Valentão, P. (2009). Fatty acid composition of wild edible mushrooms species, A comparative study. *Microchemical Journal*, 93, 29.
- [7] Öztürk, M., Tel-Cayan, G., Muhammad, A., Terzioglu, P., & Duru, M. E. (2015). Mushrooms, a source of exciting bioactive compounds, In, Atta-ur-Rahman, editor. *Studies in Natural Product Chemistry*. Amsterdam: Elsevier, 45, 363.
- [8] Duru, M. E., & Tel-Çayan, G. (2015). Biologically active terpenoids from mushroom origin, A Review. *Records of Natural Products*, 9, 456.
- [9] Zhang, G., Sun, J., Wang, H. ve Ng, T.B. (2010) First isolation and characterization of a novel lectin with potent antitumor activity from a *Russula* mushroom, *Phytomedicine*, 17-775-781.
- [10] Diez, V. A., & Alvarez, A. (2001). Compositional and nutritional studies on two wild edible mushrooms from northwest Spain. *Food Chemistry*, 75, 417.
- [11] Agrahar-Murugkar D, Subbulakshmi G. Nutritional value of edible wild mushrooms collected from the Khasi hills of Meghalaya. *Food Chemistry* 89, 599–603, 2005.
- [12] Manzi P, Aguzzi A, Pizzoferrato L. Nutritional value of mushrooms widely consumed in Italy. *Food Chemistry* 73, 321–325, 2001.
- [13] Sanmee, R, Dell B, Lumyong P, Izumori K, Lumyong S. Nutritive value of popular wild edible mushrooms from northern Thailand. *Food Chemistry* 84, 527–532, 2003.
- [14] Yilmaz, N., Türkekul, I., Bulut, S., Sahin, F., & Bayrak, Ö. F. (2013). Fatty Acid Composition in Ten Mushroom Species Collected from Middle Black Sea Region of Turkey. *Asian Journal of Chemistry*, 25, 1216.
- [15] Sarikahya, N. B., Ucar, E. O., Kayce, P., Gokturk, R. S., Sumbul, H., Arda, N., & Kirmizigil, S. (2015). Fatty Acid Composition and Antioxidant Potential of Ten Cephalaria Species. *Records of Natural Products*, 9, 116.
- [16] Tel, G., Deveci, E., Küçükaydın, S., Özler, M. A., Duru, M. E., & Harmandar, M. (2013). Evaluation of Antioxidant Activity of *Armillaria tabescens*, *Leucopaxillus gentianeus* and *Suillus granulatus*, the mushroom Species from Anatolia. *Eurasian Journal of Analytical Chemistry*, 8, 136.
- [17] Türkoğlu, A., Duru, M. E., & Mercan, N. (2007). Antioxidant and Antimicrobial Activity of *Russula delica* Fr, An Edible Wild Mushroom. *Eurasian Journal of Analytical Chemistry*, 2, 1.
- [18] Olennikov, D. N., Agafonova, S. V., Penzina, T. A., & Borovskii, G. B. (2014). Fatty Acid Composition of Fourteen Wood-decaying Basidiomycete Species Growing in Permafrost Conditions. *Records of Natural Products*, 8, 184.

- [19] Mau, J.L., Chen, P.R., Yang J.H. (1998). Ultraviolet Irradiation Increased Vitamin D2 Content in Edible Mushrooms. *J. Agric. Food Chem*, 46, 5269–5272.
- [20] Yazawa, Y., Yokota, M., Sugiyama, K. (2000). Antitumor promoting effect of an active component of Polyporus, ergosterol and related compounds on rat urinary bladder carcinogenesis in a short-term test with concanavalin A. *Biol Pharm Bull*. Nov;23(11):1298-302.
- [21] Shao, SQ., Hernandez, M., Kramer, JKG., Rinker, DL., Tsao, R. (2010) Ergosterol profiles, fatty acid composition, and antioxidant activities of button mushrooms as affected by tissue part and developmental stage. *J Agr Food Chem* 58:11616-11625.
- [22] Kuo CF, Hsieh CH, Lin WY (2011) Proteomic response of LAP-activated RAW 264.7 macrophages to the anti-inflammatory property of fungal ergosterol. *Food Chem* 126:207-212.
- [23] Hu SH, Liang ZC, Chia YC, et al. (2006) Antihyperlipidemic and antioxidant effects of extracts from *Pleurotus citrinopileatus*. *J Agr Food Chem* 54:2103-2110.
- [24] James A. Duke (2008), *Storax (Liquidambar orientalis Mill. and L., Styraciflua L.)*", *Duke's Handbook of Medicinal Plants of the Bible, Taylor & Francis*, pp. 258–259.
- [25] Öztürk, M., Çelik, A., Güvensen, A., Hamzaoğlu, E. (2008). Ecology of tertiary relict endemic *Liquidambar orientalis* Mill. Forests. *Forest Ecology and Management*, Volume 256, Issue 4, Pages 510–518.