

***Thymbra Spicata* Var. *Intricata* Induces Mesenchymal Stem Cell Proliferation and Osteogenic Differentiation**

Aysegül Mendi^{1*}, Beyza Gökçınar Yağcı² Mustafa Kızıloğlu³, Nurdan Saraç⁴, Aysel Uğur¹, Derviş Yılmaz³, Duygu Uçkan².

¹ Gazi University, Faculty of Dentistry, Department of Medical Microbiology, Ankara, Turkey; ² Hacettepe University, PEDI-STEM Center for Stem Cell Research and Development, Ankara, Turkey; ³ Gazi University, Faculty of Dentistry Department of Oral and Maxillofacial Surgery, Ankara, Turkey; ⁴ Mugla Sıtkı Kocman University, Faculty of Sciences, Department of Biology, Mugla.

ABSTRACT

*A natural agent that maintains mesenchymal stem cell (MSCs) viability, promotes osteogenic differentiation while modulating the immunological response could achieve success in regeneration during healing and may also prevent bone resorption and improve regeneration. We aimed to demonstrate that a *Thymbra spicata* var. *intricata* extract could induce proliferation, differentiation, and modulate the immune responses of mesenchymal stem cells (MSCs). Using xCELLigence, a real-time monitoring system, we obtained a growth curve of MSCs. A dose of 10 µg/mL was the most efficient concentration for vitality. Osteogenic differentiation and antiinflammatory activities were determined by using an ELISA Kit to detect early and late markers of differentiation. The Osteonectin (ON, early osteogenic marker) level decreased while the Osteocalcin (OCN, late osteogenic marker) level increased in the *T. spicata* var. *intricata* treated group, suggesting that *T. spicata* var. *intricata* may accelerate osteogenic differentiation. Reduced level of the IL-6 cytokine in response to TNF-α was evident. *T. spicata* var. *intricata* could be a promising osteogenic inducer in dentistry and could be used safely in biocomposites or scaffold fabrications.*

Key words: mesenchymal stem cell, osteogenic differentiation, *T. spicata* var. *intricata*, bone regeneration

* Author for correspondence: aysegulmendi@gmail.com

INTRODUCTION

Dental pulp mesenchymal stem cells (DP-MSCs) are a type of mesenchymal stem cell (MSCs) found in the cell-rich zone of the pulp tissue of teeth¹. DP-MSCs have a strong self-renewal ability and the potential for multi-directional differentiation, which gives them great therapeutic potential for repairing damaged and/or defective tissue².

The presence and maintenance of alveolar bone is tooth dependent. After tooth extraction, the alveolar bone is slowly resorbed down to the body of the jaw bones. In cases of complete tooth loss, there is progressive bone resorption, which can result in extensive atrophy of the jaw bones and lead to major clinical challenges for implant placement and the construction of dental prostheses³. Even without therapeutic intervention, the periodontium can exhibit a significant capacity for regeneration. However, such endogenous activity has limited capacity for periodontal regeneration. In addition, the impairment of bone formation increases in patients with osteoporosis and diabetes mellitus and related conditions⁴. We hypothesize that a natural agent that maintains MSCs viability, promotes osteogenic differentiation while modulating the immunological response could achieve success in regeneration during healing and may also prevent bone resorption and improve regeneration.

Thymbra spicata L. a wild medicinal plant of Eastern Mediterranean countries; is represented by namely *T. spicata* var. *spicata* and *T. spicata* var. *intricata* in Turkey⁵. The main component, Carvacrol, predominates (>80%) in the essential oil and demonstrated significant level of anti-hypercholesterolaemic and antioxidant activities in mice⁶. Also carvacrol has been observed with many diverse physiological actions, such as antibacterial⁷, antiinflammatory⁸. Dried herbal parts of the plant are used as herbal tea, and in folk medicine to treat asthma, colic, bronchitis, coughs, diarrhoea and rheumatism in various parts of Turkey⁹. It was important for our study that dried *T. spicata* herb contain 1925.7 mg/kg Ca since experimental evidence clearly indicates the key role of Ca²⁺ in osteoinduction^{10, 11}.

Although various physiological activities of *T. spicata* have been demonstrated, its link to osteogenic differentiation of mesenchymal stem cells has never been explored. In the present study, we hypothesized that *T. spicata* var. *intricata* could maintain the viability of DP-MSCs, induce promote their osteogenic differentiation, which may enable the successful regeneration of hard tissues. We also used bone marrow derived MSCs as control.

MATERIALS AND METHODS

Extraction of Plant Samples

T. spicata var. *intricata* flower buds were purchased from the local market in Mugla, Turkey. The air-dried plant samples were extracted with ethanol (Merck, Taufkirchen Germany) using a Soxhlet apparatus. The extracts were evaporated and stored in sterile opaque glass bottles under refrigerated conditions until use. The dried extract was prepared in DMEM-LG with 10% foetal bovine serum (FBS) (Invitrogen, Carlsbad, Calif., USA), 1% L-glutamine (Sigma, Taufkirchen Germany) and 1% Penicillin-Streptomycin (Invitrogen, Carlsbad, Calif., USA) for the studies.

Isolation and Culture of Dental Pulp Mesenchymal Stem Cells

Human dental pulp tissue was obtained from patients (15-20 years of age) who were undergoing extraction of their third molars for orthodontic reasons at the Department of Oral and Maxillofacial Surgery, University of Gazi, Ankara. All patients provided

informed consent (Ethics Commit. Rep. No: G.Ü. B30.2. GÜN 0.21.71.00). After the tooth surfaces were disinfected, the teeth were mechanically fractured, and the dental pulp was gently isolated with forceps. The pulp tissue was rinsed in α -MEM supplemented with 2 nM L-glutamine, 100 U/mL penicillin, 100 μ g/mL streptomycin and 10% fetal bovine serum (FBS, Invitrogen) (hereafter referred to as the MSC culture medium), after which it was minced into fragments of 1 to 2 mm³. The tissue fragments were cultured in T75 Nunc plates in the MSC culture medium at 37°C in a humidified atmosphere containing 5% CO₂. Human BM-MSCs were a kind gift from Hacettepe University Center for Pediatric Stem Cell and Research and Development. BM-MSCs were suspended at a concentration of 1x10⁶ cells/mL in MSC culture medium. The culture media was changed every 2 to 3 days, and the cell cultures were monitored regularly with an inverted microscope (Olympus CKX41, Tokyo, Japan). Upon reaching 70-80% confluence, the cells were harvested with 0.05% Trypsin/EDTA (Sigma, Taufkirchen Germany) and sub-cultured for further experiments.

Immunophenotypic Analysis

The culture-expanded adherent cells were analysed by flow cytometry (BD FACSAria, USA). The antibody panel included CD29- FITC (e-bioscience, USA); CD73-PE (BD, USA), CD 90-PE (BD, USA), CD44-PE (e-bioscience, USA) as mesenchymal stromal markers, as well as their isotype controls. CD45-FITC (BD, USA); CD14-PE (BD, USA); and CD34-FITC (BD USA) were used as haematopoietic markers to exclude cells of haematopoietic origin. The relative frequencies of the cells that expressed the respective surface markers were analysed using FACS Diva software 6.0.0 (BD) by acquiring 10,000 events for each sample.

Effect Of *T. spicata* var. *intricata* on Proliferation of the MSCs Using the xCELLigence System

Initially we examined the proliferation of DP-MSCs in a 24 well culture microplate seeded at a density of 5000 cell/cm². DP-MSCs were cultured with different concentrations (1, 3, 5, 10, 25, 50, 75, and 100 μ g/mL) of *T. spicata* var. *intricata* up to the control group had 90% confluency. Cells were counted by trypan blue method and the three concentrations which induced the cell number was selected for xCELLigence analysis (data not shown). The xCELLigence system was used according to the manufacturer's instructions¹². Briefly, the E-plate 96 was connected to the xCELLigence system and verified in the cell culture incubator to ensure that proper electrical contacts were established, and the background impedance was measured. Subsequently 100 μ l of MSCs culture media containing 5, 10 and 25 μ g/mL *T. spicata* var. *intricata* extract and standart culture media as control were added into each well of E-plate 96. Meanwhile, the cells were resuspended (5000 cells/cm²) in MSC culture media for their concentration. 100 μ l of each cell suspension was added to each well, in order to determine effect of *T. spicata* var. *intricata* extract on cell proliferation. Cell growth and proliferation were monitored every 30 min for up to 290 h. The growth curve, cell index, and doubling time (DT) values were determined.

Effect of *T.spicata* var. *intricata* on MSCs Differentiation

The concentration that decreased the doubling time and increased the proliferation was selected based on the results from the xCELLigence system analysis. The selected concentration was added to the osteogenic and adipogenic differentiation media¹³. The images were obtained with a CKX41 digital imaging microscope (Olympus, Tokyo, Japan). The secreted Osteocalcin (OCN) and Osteonectin (ON) levels in the supernatants were assessed using an ELISA kit according to the

manufacturer's instructions (R&D Systems, Inc. Minneapolis). The limits of detection for the ELISA were 1.2 to 75 ng/mL for OCN and 1.56 to 50 ng/mL for ON.

Determining the Immunomodulatory Activities

DP- and BM-MSCs were plated at a density of 5000 cell/cm² in 96-well culture plates and allowed to attach overnight. The cells were pretreated with 10 µg/mL *T. spicata* var. *intricata* extract for 1 h, and 10 ng/mL TNF-α were then added. After 24 h, the cell culture supernatants were collected and stored at -80°C for use in the IL-6, IL-10 ELISAs, according to the manufacturer's instructions. The ELISA limits were 0,052-0,118 pg/mL for IL-6 and 0,39-25 pg/mL for IL-10. Media alone, with TNF-α, and with *T. spicata* var. *intricata* were included as controls.

Statistical Analysis

All calculations were performed using the RTCA integrated software of the xCELLigence system. The RTCA software fits the curve of the selected sigmoidal dose response equations to the experimental data points. The data are presented as the mean (µg/mL) ± SD (n=4). For the proliferation experiments, the statistical analysis was performed using one-way analysis of variance (ANOVA) ($p < 0.05$).

RESULTS

Identification of MSCs

The common MSC markers (CD29, CD73, CD44, and CD90) were constitutively positive (>95%) and the hematopoietic markers (CD14, CD34, and CD45) were negative (>95) in all samples tested, indicating a mesenchymal origin of the cells (Figure 1).




Figure 1: Surface markers of DP-MSCs and BM-MSCs

xCELLigence Assays

Trypan blue assay showed that the concentrations between 1-5 µg/mL and 25-100 µg/mL reduced the viability of DP-MSCs. 10µg/mL was found close to the Control group (data not shown). We selected 5, 10, and 25 µg/mL for the xCELLigence analysis system (Figure 2). The xCELLigence analysis showed that the growth

curves of the DP-MSCs and BM-MSCs treated with *T. spicata* var. *intricata* were slightly lower than the growth curve of the controls, but they were close to each other for the culture period except 5 $\mu\text{g/mL}$. We determined the cell indexes for the time intervals of the lag, log, and plateau phases using the growth curve. xCelligence assay showed that cells treated with *T. spicata* var. *intricata* had lower adhesion, but similar viability when compared with control cells. The doubling time (DT) was reduced at 10 $\mu\text{g/mL}$ in DP- and BM-MSCs. Thus, *T. spicata* var. *intricata* could be a good proliferation inducer. The IC_{50} value was found 7,7 $\mu\text{g/mL}$ for DP-MSCs and 17 $\mu\text{g/mL}$ for BM-MSCs at 290 h.




Figure 2: xCELLigence analysis of DP-MSCs and BM-MSCs. Growth curve of the cells. a1) DP-MSCs, b1) BM-MSCs; doubling time of the DP-MSCs (a2) and BM-MSCs (b2) and the cell indexes at lag mid log, and stationary phases (c) of the cells were determined.

Differentiation Assays of MSCs

Subsequently, the characteristics features of the cells were studied. (Figure 3). Adipogenic differentiation exhibited a quite difference in DP-MSCs treated with *T. spicata* var. *intricata*. Approximately 20% of the cells became rounder; however, no lipid droplets were observed in DP-MSCs. In contrast to adipogenic differentiation, the DP-MSCs underwent rapid osteogenic differentiation. Calcium granules similar to bone nodules were seen in BM-MSCs. *T. spicata* var. *intricata* treated DP- and BM-MSCs showed more osteogenic differentiation than the control cells. We also determined the levels of the ON and OCN osteogenic markers.




Figure 3: Differentiation potential of DP-MSCS (I) and BM-MSCS (II). adipogenic differentiation was not shown in DP-MSCs (I.a1) while BM-MSCs (II.a1) were well differentiated. Although *T. spicata* var. *intricata* lead to cell morphology changes in adipogenic differentiation of DP-MSCs (I.b1), it was not occurred like BM-MSCs (II.b1). Osteogenic differentiation was observed in DP-MSCs (I.a2) and BM-MSCs (II.b2). The extract of *T. spicata* var. *intricata* induced both DP-MSCs (I.b2) and BM-MSCs (II.b2) osteogenic differentiation (4x, Olympus CKX41, Japan). Osteonectin (III) and osteocalcin (IV) levels are calculated from supernatant of culture medium.

Determining the Preventive Effect of *T. spicata* var. *intricata* on the Inflammatory Response of MSCs Following TNF- α Stimulation

Figure 4 shows the antiinflammatory activities of *T. spicata* var. *intricata*. Obtained data showed an antinflammatory and immunomodulatory effect of the extract.




Figure 4: IL-6 (a) and IL-10 (b) level determined in the culture supernatant of DP and BM-MSCs.

DISCUSSION

In the present study, we demonstrated that *T. spicata* var. *intricata* extract can induce proliferation, differentiation, and modulate the immune responses of DP- and BM-MSCs. Initially, we aimed to define the effective concentration of *T. spicata* var. *intricata*. Using the xCELLigence system, we obtained a growth curve for MSCs

which provides information on three parameters: the lag phase before cell proliferation is initiated after subculture, the DT in the middle of the exponential growth phase, and the terminal density². The lag phase is the time period when the cells are recovering from trypsinization, synthesizing new extracellular matrix and cytoskeleton, and adapting for attachment, spreading and re-entering the cell cycle. Cells do not divide in the lag phase¹³. We could conclude that *T. spicata* var. *intricata* reduced the adhesion of DP-MSCs at 48h and 98h whilst BM-MSCs were adhered and proliferated more than the Control group. Ideally, towards the end of the log phase, the culture becomes confluent and its growth rate reduces, and in some cases, cell proliferation ceases. At this stage, the culture enters the plateau, or stationary phase, in which cell division is balanced by cell loss. DP-MSCs were still alive where BM-MSCs began to reduced their cell number. Also *T. spicata* var. *intricata* reduced the DT of DP- and BM-MSCs suggesting that it would be an advantage in if used as a filling material scaffold in dental therapies.

Oriental medicine practices are primarily based on personal experience, which often result in unknown mechanisms and difficulties in dose specification. We used a real-time monitored system, xCELLigence, to determine the effective concentration. The xCELLigence system is much more sophisticated than other conventional endpoint, cell-based assays. Real-time and continuous monitoring enable the label-free assessment of cell proliferation, viability, and cytotoxicity by showing the physiologic state of the cells and eliminating expensive reagents that are used in conventional cell analyses¹⁴.

The osteogenic differentiation potentials of DP-MSCs in vitro and in vivo have been well documented in a variety of studies^{2, 15}. Adipogenic differentiation was not seen in DP-MSCs while BM-MSCs were well differentiated. Our findings were agree with those of Gronthos et al², who expanded DP-MSCs from single-cell clones and demonstrated that they exhibited osteogenic differentiation and did not form lipid-laden adipocytes. During osteogenic cell differentiation, the markers of the undifferentiated cells are gradually turned off, and the differentiation markers are sequentially expressed. We observed the sequential secretion of proteins at the end of the assay, in which the ON levels decreased in the *T. spicata* var. *intricata*-treated group compared to the control group. ON is an early marker of osteogenesis that is synthesized by preosteoblasts and has less affinity to collagen. The ON transcript is quite stable, with a half-life of >24 hours under conditions of transcription arrest¹⁶. The *T. spicata* var. *intricata*-treated DP- and BM-MSCs exhibited higher OCN levels than the untreated DP- and BM-MSCs. We suggest that *T. spicata* var. *intricata* accelerates the differentiation of MSCs. Therefore, *T. spicata* var. *intricata* could be a safe inducer for both healthy and medically compromised patients.

Both IL-6 and IL-10 were present in the DP- and BM-MSCs cell culture supernatants⁴. Our results showed that when the extract used alone the IL-6 level was increased in BM-MSCs while it was decreased in DP-MSCs. On the other hand in contrast with this result, the DP-MSCs were pretreated with *T. spicata* var. *intricata*, before TNF- α stimulation the IL-6 level decreased at a ratio of 94% and 42% in DP-MSCs and BM-MSCs respectively. This result raise the possibility that the number of IL-6 receptor on cell membrane may be important factor controlling the IL-6 response in target tissues in physiological or pathological conditions. In similar with our results some reports suggested that when soluble IL-6 receptor is available, IL-6 additionally acts on osteoblasts to induce their differentiation¹⁷. Increased osteogenic activity could be linked to the induced IL-6 activity in BM-MSCs. IL-6 is a multifunctional cytokine that regulates pleiotropic functions of cells and tissues¹⁸. Reduced IL-6 in response to TNF- α showed that *T. spicata* var. *intricata* could be a well immunomodulatory agent in inflammatory conditions. There have been many reports which suggest that IL-6 plays an important role in osteoclastic bone

resorption in vitro as well as in vivo¹⁹. According to our results and the reported data a coculture study with osteoblasts and osteoclast treated with *T. spicata* var. *intricata* should be done.

Here we demonstrated a *T. spicata* var. *intricata* extract can induce osteogenic differentiation, and modulate the immune responses of DP-MSCs and BM-MSCs. To introduce a new material to be used as a new scaffold in bone regeneration cell growth properties and osteogenic stimulation potency should be determined, and in vitro cell culture studies are required to carry the new material to the stage of animal studies. *T. spicata* var. *intricata* could be a promising osteogenic inducer in dentistry and could be used safely in biocomposites or scaffold fabrications.

ACKNOWLEDGEMENT

This study was supported by the Turkish Scientific and Technological Research Council (TUBITAK), Project no: SBAG 113S448. We specially thank to Prof. Dr. Petek Korkusuz and Sevil Köse for assistance in xCELLigence assays and analysis.

CONFLICT OF INTERESTS

Authors declare that there is no conflict of interests.

REFERENCES

1. Estrela C, Alencar AHGD, Kitten GT, Vencio EF, Gava E. Mesenchymal stem cells in the dental tissues: perspectives for tissue regeneration. *Braz Dent J.* 2011; 22: 91–8.
2. Gronthos S, Mankani M, Brahım J, Robey PG, Shi S. Postnatal human dental pulp stem cells (DPSCs) in vitro and in vivo. *Proceed Nat Acad Sci.* 2000; 97: 13625–30.
3. Hoiruchi K, Amizuka N, Takeshita S, Takamatsu H, Katsuura M, Ozawa H, Toyama Y, Bonewald LF, Kudo A. Identification and characterization of a novel protein, perostin, with restricted expression to periosteum and periodontal ligament and increased expression by transforming growth factor beta. *J Bone Miner Res.* 1999; 14: 1239–49.
4. Egermann M, Goldhahn J, Schneider E. Animal models for fracture treatment in osteoporosis. *Osteoporos Int.* 2005;16 (suppl2): 129–38.
5. Tanker M, Ilisulu F. *Thymbra spicata* L. var. *spicata*: one of the plants used in Turkey as thyme. *Turk J Bot.* 1984; 8: 104–108.
6. Akkol EK, Avci G, Kucukkurt I, Keles H, Tamer U, Ince S, Yeşilada, E. Cholesterol-reducer, antioxidant and liver protective effects of *Thymbra spicata* L. var. *spicata*. *J Ethnopharmacol.* 2009;126: 314–9.
7. Chami F, Chami N, Bennis S, Trouillas J, Remmal A. Evaluation of carvacrol and eugenol as prophylaxis and treatment of vaginal candidiasis in an immunosuppressed rat model. *J Antimicrob Chemo.* 2004; 54: 909–14.
8. Landa P, Kokoska L, Pribylova M, Vanek T, Marsik P. In vitro anti-inflammatory activity of carvacrol: inhibitory effect on COX-2 catalyzed prostaglandin E(2) biosynthesis. *Arch Pharm Res.* 2009; 32: 75–8.
9. Baser KHC. Her Derde Deva Bir Bitki-Kekik. *Bilim ve Teknik.* 2001;74: 4
10. Kizil S. Determination of essential oil variations of *Thymbra spicata* var. *spicata* L. naturally growing in the wild flora of East Mediterranean and Southeastern Anatolia regions of Turkey. *Indust Crops and Prod.* 2010;32: 593-600.
11. Chai YC, Roberts SJ, Schrooten J, Luyten FP. Probing the osteoinductive effect of calcium phosphate by using an in vitro biomimetic model. *Tiss Eng Part A.* 2011; 17: 1083–97.
12. Roche Diagnostics. GmbH. Introduction of the RTCA SP Instrument. RTCA SP Instrument Operator's Manual, A.Acea Biosciences, Inc ; 2008; 14–16.
13. Pittenger MF, Mackay AM, Beck S. Multilineage potential of adult human mesenchymal stem cells. *Sci.* 1999; 284: 1168–70.
14. Hensten-Pettersen A. Comparison of the methods available for assessing cytotoxicity. *Int Endod J.* 1988;21: 89-99.

15. D'aquino R, Graziano A, Sampaolosi M, Laino G, Pirozzi G, De Rosa A, Papaccio G. Human postnatal dental pulp cells co-differentiate into osteoblasts and endotheliocytes: a pivotal synergy leading to adult bone tissue formation. *Cell Death Differ.* 2007; 14: 1162–71.
16. Dole NS, Kapinas K, Kessler CB, Yee S, Adams JD, Pereira RC, Delany AM. A single nucleotide polymorphism in osteonectin 3' untranslated region regulates bone volume and is targeted by miR-433. *J Bone and Mineral Res.* 2015; 30: 723–32.
17. Ishihara K, Hirano T. IL-6 in autoimmune disease and chronic inflammatory proliferative disease. *Cytokine Growth Factor Rev.* 2002; 13:357–68.
18. Taga T, Kishimoto T. Role of a two-chain IL-6 receptor system in immune and hematopoietic cell. *Crit Rev Immunol.* 1992;11:265-80.
19. Roodman, GD. Interleukin 6: an osteotropic factor?. *J. Bone Miner Res.* 1992;7:475-8.

Received: February 03, 2016;
Accepted: July 14, 2016