

Research

Return to work in miners following anterior cruciate ligament reconstruction

Ugur Tiftikci¹, Sancar Serbest^{1,*}, Cem Yalin Kilinc², Gül Öznur Karabicak³, Özge Vergili⁴

¹Kirikkale University, Faculty of Medicine, Department of Orthopaedics and Traumatology, Kirikkale, Turkey, ²Mugla University, Faculty of Medicine, Department of Orthopaedics and Traumatology, Mugla, Turkey, ³Hacettepe University, Faculty of Health Sciences, Department of Physical Therapy and Rehabilitation, Ankara, Turkey, ⁴Kirikkale University, Faculty of Health Sciences, Department of Physical Therapy and Rehabilitation, Kirikkale, Turkey

*Corresponding author: Sancar Serbest, Kirikkale University, Faculty of Medicine, Department of Orthopaedics and Traumatology, Kirikkale, Turkey

Key words: Anterior cruciate ligament, injury, knee, reconstruction, miner, return to work

Received: 15/09/2015 - Accepted: 17/10/2015 - Published: 22/10/2015

Abstract

Introduction: The aim of the study is retrospectively investigated durations for returning to work following anatomic ACL reconstruction by hamstring autograft in miners and the reasons in patients who were delayed to return to work. **Methods:** Miners with symptomatic anterior cruciate ligament rupture underwent arthroscopic reconstruction. Patients were evaluated in terms of range of motion (ROM) values; Lysholm, Cincinnati and Tegner activity scales; laxity testing and complications. By modifying the method used by Fitzgerald et al. we decided for the criteria returning to work. **Results:** Thirty three patients were evaluated with mean followup of 22.7 ± 8.3 months (range 13-46 months). Mean age at the surgery was 27.8 (18-38) years. Lysholm, Cincinnati and Tegner activity scales were significantly higher from preoperative scores (Lysholm scores: preoperative: 60.7 ± 12.5 , postoperative: 90.3 ± 4.8 ($P < 0.001$); Tegner activity scores: Preoperative 3.5 ± 1.4 , postoperative: 6.2 ± 1.5 ($P < 0.001$); Cincinnati scores: Preoperative: 14.8 ± 5.3 , postoperative: 26.9 ± 1.6 ($P < 0.001$). The average time for returning to work was determined as 15.3 ± 4 weeks. There was no significant difference for knee scores and time for returning to work between patients with meniscal injuries and don't have meniscus lesions. **Conclusion:** The reasons for delays in returning to work was work accident. Hematoma or effusion and pain inside the knee were the most significant reason which affected returning to work.

Pan African Medical Journal. 2015; 22:173 doi:10.11604/pamj.2015.22.173.7979

This article is available online at: <http://www.panafrican-med-journal.com/content/article/22/173/full/>

© Ugur Tiftikci et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Anterior cruciate ligament (ACL) injury is the most commonly injury among ligament injuries of the knee region. ACL reconstruction surgery is the most frequently performed operation by the orthopedic surgeons [1]. This injury has been shown to commonly cause joint effusion, muscle weakness, loss of joint function, joint instability, pain and osteoarthritis at young ages, between 10-20 years of age [2-4]. The management of ACL injuries is both conservative and surgical. However, the up-to-date management has increased the tendency to surgical treatment in people dealing with sports actively, young people, active people and heavy-workers [1,2,5]. The most preferred and up-to-date method in surgical treatment is "medial portal anatomic ACL reconstruction by hamstring autograft" [5]. In this study, we retrospectively investigated durations for returning to work following anatomic ACL reconstruction by hamstring autograft in miners and the reasons in patients who were delayed to return to work.

Methods

33 of 38 miners who had undergone arthroscopic ACL reconstruction by using anatomic medial portal between 2009 and 2014 were included in the study, through their hospital records. Patients with multiple ligament injuries, total meniscectomy, contralateral knee or lower extremity injury, who were aged over 40 years, and whose records were inaccessible were excluded from the study. Laxity tests (Anterior drawer, Lachmann, Pivot shift), effusion and tenderness were assessed. ROM value was scored using goniometer and differences in maximum ROM between involved and contralateral sides were measured. Lysholm, Cincinnati and Tegner activity scale was scored after 6 months of the surgery. However, since KT-1000 device was absent, arthrometer test was not performed. The time for starting to work and the reasons for delay in returning to work were identified through hospital records. By modifying the method used by Fitzgerald et al. [6], we decided for the criteria returning to work by knee function test, single-legged hop test score, Lachman test, anterior drawer test and pivot-shift test in miners. This study was approved by the institutional ethical committee; all patients were informed and the study was performed in accordance with the declaration of Helsinki as revised in 2000.

Surgical Technique

1 gr of first-generation cephalosporin was administered to all patients half an hour before the operation for prophylaxis. This prophylactic antibiotic was continued for 24 hours, until the drain was removed. For confirmation of the ACL tear, in all patients, before surgery, the arthroscopic assessment of the knee was performed in the supine position under general or regional anesthesia and under pneumatic tourniquet. The associated pathologies were identified and necessary treatments were made arthroscopically. By using a 4 cm longitudinal incision, at the medial of tibial tubercle and over pes anserinus, gracilis and semitendinosus tendons were reached. By using a tendon stripper, tendons were excised and the attached muscle tissues were cleaned. By folding the tendons four times, the diameter of the graft was determined. From the appropriate medial portal for this diameter, anatomic femoral tunnel and then the tibial tunnel were prepared; then, by the help of guidewires, graft was positioned into the tunnels. At the proximal side, the graft was fixed to the femur by endobutton system (ToggleLoc ZipLoop, Biomet Sport Medicine). Distally, while the knee was positioned in 10°-20° flexion and graft was being held tightly, the fixation of the graft inside the tunnel was accomplished by one BioComposite Interference Screw (Biomet Sports Medicine) and one staple U nail.

Postoperative management protocol

Isometric quadriceps strengthening and flexion exercises were immediately started in all patients. Bearing weight as much as they can tolerate, the patients were permitted to walk using knee braces and crutches. At the 2-14 days our goals were obtained full extension, minimize swelling, allow wound healing, maintain active quadriceps control and achieve 90° of flexion. At the 2-6 weeks, we aimed to increase flexion to 135 degrees and increase muscle tone. Daily activities were allowed to be gradually increased until the 4th week. In all patients, compelling early rehabilitation was started [7,8]. In cases of muscular atrophy and functional disability of the knee, advanced physical rehabilitation therapy was used. Sports and active exercises were permitted after postoperative 2 ½ months.

Statistical analysis

Statistical analysis was carried out by using the Statistical Package for the Social Sciences (SPSS) software version 15.0 (SPSS Inc., Chicago, USA) and data are presented as mean ± standard

deviation comparisons paired t-test, unpaired t-test and Mann-Whitney U-test was performed. $P < 0.05$ was considered to be significant.

Results

The average age of patients was 27.8 (18-38). All patients were male. The reasons for ACL injury was work-accident in 16, sports injury and other reasons in 17 patients. The ACL tear was at the right knee in 20 patients and left knee in 13 patients. In the postoperative period Lysholm, Cincinnati and Tegner activity scores were significantly higher from preoperative knee scores (**Table 1**). In 12 patients, medial, in 7 patients lateral, and in 6 patients both medial and lateral meniscal tears were present. In 3 patients having meniscal injuries, partial meniscectomy was performed. In 16 patients, meniscus was sutured using all in site technique (Sequent, Linvatec. In 3 patients, medial parapatellar plica excision was performed. The mean ROM deficit (involved vs. contra knee) was significant difference for knee scores, ROM deficits and time for returning to work of the patients with work-accident and sports injury (**Table 2**). The average time for returning to work was determined as $15,3 \pm 4$ (range 10-27) weeks. The reasons for delays in returning to work was work accident. Hematoma or effusion, pain inside the knee were the most significant reason which affected returning to work (**Table 2**). During the last follow-up, the difference of muscular atrophy was found as 4.2 cm in average. There was no significant difference for knee scores and time for returning to work between patients with meniscal injuries and don't have meniscus lesions (**Table 3**). In 4 patients, for developing intra-articular effusion during the procedure, effusion drainage was performed. In one patient, at postoperative 1st month, stable U nail was removed because of superficial infection.

Discussion

ACL injuries may be managed both conservatively and surgically. In patients who sustain a sedentary life, activity modification, muscle strengthening exercises, using braces and avoiding activities which need performance are preferred in the conservative management of ACL injuries [9,10]. In these patients, the risk of developing osteoarthritis is increased by 50% in the next 10-20 years. Pain and loss of function occurs as a result of osteoarthritis [4]. However,

ACL management has progressed toward surgical treatment in the last 10 years. The reasons for this are that the patients are young and active and they want to gain back their pre-injury activity levels and knee functions in the early period [11]. The patients included in this study were working underground, in the coal mines, in harsh conditions and usually on slippery surfaces. Additionally, because of the high collapse risk and mine-transportation vehicles, they were exposed to direct trauma. For patients, returning to active work early was important in the aspect of financial losses and continuation of their business contracts. Hence, in patients with poor knee stability (Lachman, pivot-shift +), conservative therapy was not recommended.

The most commonly and up-to-date method for ACL repair is the medial portal anatomical ACL reconstruction by hamstring autograft [5]. The rapid return of patients to their pre-injury activities and endeavors has gained importance. For this reason, the postoperative rehabilitation protocols have improved significantly and compelling early rehabilitation has started to gain more acceptance [7,8]. Rapid return to sports is suggested especially within the postoperative 6 months following ACL reconstruction [12,13]. Since our patients were young, active and working in harsh conditions, we recommended surgical treatment to them, also. We rapidly initiated the compelling early postoperative rehabilitation. We did not allow our patients to perform active sportive exercises for two months. The most important reasons for delayed returning to work in patients who had undergone ACL reconstruction for work-related accidents are the severity of the trauma and psychologic reasons. In these patients, posttraumatic stress disorder is seen more frequently when compared to sports injuries. In performed studies, although many factors have been reported to contribute for delay in returning to sports in athletes, the most important one was the psychologic factor [14-17]. The second factor may be secondary gain. Since these are legally work-related accidents, these patients cannot be laid off according to their employment contracts; situations occur in which large amounts of compensation become necessary to be paid by the employer. Atrophy of the quadriceps muscle is observed in most of the patients who undergo ACL reconstruction. In the literature, the most common complications have been reported to be muscular atrophy and weakness [18,19] and insufficiencies in the strength of hamstring and quadriceps muscles, motor coordination and proprioception may continue in patients until the end of the 1st postoperative year [20].

The main reason for quadriceps weakness is not donor site morbidity, but the deterioration of the receptor and neuromuscular activation systems [21,22]. Mendias et al. in their study, determined that myostatin and TGF- β levels were elevated in patients having muscular atrophy. Future studies may be conducted on myostatin [23]. In the study conducted by De Jong et al., they determined significant strength reduction in quadriceps muscle after ACL reconstruction with hamstring tendon graft. This reduction was being gradually less within 6-12 months postoperatively; however, when compared to the preoperative period, the insufficiency still continued [24]. Muscular atrophy was observed in 75% of our patients, and this was the main cause for delay in returning to work. This patient group also required physical rehabilitation therapy.

Physical rehabilitation plays an important role for achieving successful clinical outcome, following anterior cruciate ligament (ACL) reconstruction. To return to pre-injury activity level and sports, modern treatment methods and rehabilitation protocols have been widely realized and easily accessible [25]. Although variations exist among physical rehabilitation programs, accelerated programs are preferred to promote early return to sports and active living, which involve early initiation of motion, early regaining the muscle strength and recovery of knee functions [7,8,26-28]. We, also, used an accelerated physical therapy protocol in our clinic, immediately after surgery in our patients. Closed-kinetic chain exercises for strengthening quadriceps and hamstring muscle groups and exercises for joint motion range were routinely performed by all patients. Patients in whom insufficiency was thought to be present in extensor muscle groups were directed to receive professional physical rehabilitation therapy. We suggest that, in all patients who have undergone ACL reconstruction, physical rehabilitation by a professional team will increase the success rate of the operation and will accelerate the return to active life and work. In this study, the lack of a professional physical rehabilitation team to follow the patients postoperatively was a significant deficit. It might have been the reason for delayed return of patients to active life and work. In patients who had developed effusion and hematoma, return to work was delayed. In these patients, severe muscular atrophy developed and required professional physical rehabilitation therapy. The examination findings of the knee stability were normal in patients; however, since the duration for improvement of functional scores was long, they were delayed to return to work.

The association of meniscal and chondral injuries with ACL tear is frequently met and it has been shown to influence the postoperative

healing process. Early surgical treatment, particularly in the first 6 months, has been suggested [29-31]. In patients with delayed ACL reconstruction who also had meniscal and chondral lesions, these injuries have been shown to have negative impact on the results of surgical treatment [32]. In our study, we determined that meniscal and chondral lesions had not influenced the return to work in the negative direction. The reasons for this result may be performing surgery in the early period, performing meniscal repair, and the repair of bad chondral lesions by mosaicplasty in three patients. Additionally, in these patients, not returning to work for 2 ½ months may serve as a period to complete the healing process of injuries inside the knee.

Single-legged hop test is the best testing method to assess knee functions for returning to sports in patients who have undergone ACL reconstruction [33,34]. During decision-making process for returning to work in our patients, we allowed the patients who performed knee stabilization tests and single-legged hop test 75% and over to return to work. With tearing of ACL, the positional sense of the knee or proprioception is reduced [35]. In our study, evaluation of proprioception was not made; special exercise program for proprioception was not made, either. The most important limitation of our study was being retrospective. Other limitations were the lack of K-T dynamometry test, which is a knee stability test, the lack of professional physical rehabilitation for every patient and the lack of assessment and exercise program for proprioception.

Conclusion

The most important factors which influence returning to work in miners operated for ACL are hematoma-effusion inside the knee, muscular atrophy and the long duration of rehabilitation therapy process. Since patients work in harsh conditions, they do not feel themselves ready for working. Therefore, patients who undergo operations for work-accidents are unwilling to get back to work. Physical rehabilitation therapy and psychological support should be given in these patients, similar to athletes, in order to reach their pre-injury activity levels.

Competing interests

The authors declare no competing interests.

Authors' contributions

Ugur Tiftikci, research concept and design, collection and/or assembly of data, writing the article, critical revision of the article. Sancar Serbest, research concept and design, collection and/or assembly of data, writing the article. Cem Yalin Kilinc, writing the article. Özge Vergili, collection and/or assembly of data. Gül Öznur Karabıcak, research, concept and design. All authors have read and approved the final version of the manuscript.

Tables

Table 1: Knee scores

Table 2: Comparison of outcomes between work-accident and sports injury patients

Table 3: Comparison of outcomes between patients with or without meniscus lesions

References

1. Gianotti SM, Marshall SW, Hume PA, Bunt L. Incidence of anterior cruciate ligament injury and other knee ligament injuries:a national population based study. *J Sci Med Sport*. 2009;12(6):622-7. **PubMed | Google Scholar**
2. Hewett TE, Distasi SL, Myer GD. Current concepts for injury prevention in athletes after anterior cruciate ligament reconstruction. *Am J Sports Med*. 2013;41(1):216-24. **PubMed | Google Scholar**
3. Griffin LY, Agel J, Albohm MJ, Arendt EA, Dick RW, Garrett WE et al. Non contact anterior cruciate ligament injuries: risk factors and prevention strategies. *J Am Acad Orthop Surg*. 2000;8(3):141-50. **PubMed | Google Scholar**
4. Lohmader LS, Ostenberg A, Englund M, Roos H. High prevalence of knee osteoarthritis, pain and functional limitations in female soccer players twelve years after anterior cruciate ligament injury. *Arthritis Rheum*. 2004;50(10):3145-52. **PubMed | Google Scholar**
5. Middleton KK, Hamilton T, Irrgang JJ, Karlsson J, Harner CD, Fu FH. Anatomic anterior cruciate ligament (ACL) reconstruction:a global perspective; Part 1. *Knee Surg Sports Traumatol Arthrosc*. 2014;22(7):1467-82. **PubMed | Google Scholar**
6. Fitzgerald GK, Axe MJ, Snyder-Mackler L. A decision-making scheme for returning patients to high level activity with nonoperative treatment after anterior cruciate ligament rupture. *Knee Surg Sports Traumatol Arthrosc*. 2000;8(2):76-82. **PubMed | Google Scholar**
7. Majima T, Yasuda K, Tago H. Rehabilitation after hamstring anterior cruciate ligament reconstruction. *Clin Orthop Relat Res*. 2002;397:370-80. **PubMed | Google Scholar**
8. Beynon BD, Johnson RJ, Fleming BC. The science of anterior cruciate ligament rehabilitation. *Clin Orthop Relat Res*. 2002;402:9-20. **PubMed | Google Scholar**
9. Ciccotti MG, Lombardo SJ, Nonweiler B, Pink M. Non operative treatment of ruptures of the anterior cruciate ligament in middle-aged patients: results after long-term follow-up. *J Bone Joint Surg Am*. 1994;76(9):1315-21. **PubMed | Google Scholar**
10. Buss DD, Min R, Skyhar M, Galinat B, Warren RF, Wickiewicz TL. Nonoperative treatment of acute anterior cruciate ligament injuries in selected groups of patients. *Am J Sports Med*. 1995;23(2):160-5. **PubMed | Google Scholar**
11. Eitzen I, Moksnes H, Snyder-Mackler L, Engebretsen L, Risberg MA. Functional tests should be accentuated more in the decision for ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc*. 2010;18(11):1517-25. **PubMed | Google Scholar**
12. Cascio BM, Culp L, Cosgarea AJ. Return to play after anterior cruciate ligament reconstruction. *Clin Sports Med*. 2004;23(3):395-408. **PubMed | Google Scholar**

13. Reid A, Birmingham TB, Stratford PW, Alcock GK, Giffin JR. Hop testing provides a reliable and valid outcome measured during rehabilitation after anterior cruciate ligament reconstruction. *Phys Ther.* 2007; 87(3):337-49. **PubMed | Google Scholar**
14. Kvist J, Ek A, Sporrstedt K, Good L. Fear of re-injury: a hindrance for returning to sports after anterior cruciate ligament reconstruction. *Knee Surg Sport Traumatol Arthrosc.* 2005;13(5):393-7. **PubMed | Google Scholar**
15. Tripp, DA, Stanish, W, Ebel-Lam, A, Brewer, BW, Birchard, J. Fear of reinjury, negative affect, and catastrophizing predicting return to sport in recreational athletes with anterior cruciate ligament injuries at 1 year postsurgery. *Rehabil Psychol.* 2007;52:74-81. **PubMed | Google Scholar**
16. Gobbi A, Francisco R. Factors affecting return to sports after anterior cruciate ligament reconstruction with patellar and hamstring graft: a prospective clinical investigation. *Knee Surg Sport Traumatol Arthrosc.* 2006;14(10):1021-8. **PubMed | Google Scholar**
17. Langford JL, Webster KE, Feller JA. A prospective longitudinal study to assess psychological changes following anterior cruciate ligament reconstruction surgery. *Br J Sport Med.* 2009;43(5):377-88. **PubMed | Google Scholar**
18. McHugh MP, Tyler TF, Browne MG, Gleim GW, Nicholas SJ. Electromyographic predictors of residual quadriceps muscle weakness after anterior cruciate ligament reconstruction. *Am J Sports Med.* 2002;30(3):334-39. **PubMed | Google Scholar**
19. Yasuda K, Ohkoshi Y, Tanabe Y, Kaneda K. Quantitative evaluation of knee instability and muscle strength after anterior cruciate ligament reconstruction using patellar and quadriceps tendon. *Am J Sports Med.* 1992;20(4):471-75. **PubMed | Google Scholar**
20. Yosmaoglu HB, Baltaci G, Kaya D, Özer H, Atay A. Comparison of functional outcomes of two anterior cruciate ligament reconstruction methods with hamstring tendon graft. *Acta Orthop Traumatol Turc.* 2011;45(4):240-7. **PubMed | Google Scholar**
21. Krogsgaard MR, Dyhre-Poulsen P, Fischer-Rasmussen T. Cruciate ligament reflexes. *J Electromyogr Kinesiol.* 2002;12(3):177-82. **PubMed | Google Scholar**
22. Williams GN, Chmielewski T, Rudolph K, Buchanan TS, Snyder-Mackler L. Dynamic knee stability: current theory and implications for clinicians and scientists. *J Orthop Sports Phys Ther.* 2001;31(10):546-66. **PubMed | Google Scholar**
23. Mendias CL, Lynch EB, Davis ME, Sibilsky Enselman ER, Harning JA, Dewolf PD et al. Changes in Circulating Biomarkers of Muscle Atrophy, Inflammation and Cartilage Turnover in Patients Undergoing Anterior Cruciate Ligament Reconstruction and Rehabilitation. *Am J Sports Med.* 2013;41(8):1819-26. **PubMed | Google Scholar**
24. De Jong S, vanCaspel D, vanHaeff M, Saris DBF. Functional assessment and muscle strength before and after reconstruction of chronic anterior cruciate ligament lesions. *Arthroscopy.* 2007;23(1):21-8. **PubMed | Google Scholar**
25. Schoderbek RJ Jr, Treme GP, Miller MD. Bone-patellar-tendon-bone auto graft anterior cruciate ligament reconstruction. *Clin Sports Med.* 2007;26(4):525-47. **PubMed | Google Scholar**
26. Shelbourne KD, Nitz P. Accelerated rehabilitation after anterior cruciate ligament reconstruction. *Am J Sports Med.* 1990;18(3):292-9. **PubMed | Google Scholar**
27. Yabroudi MA, Irrgang JJ. Rehabilitation and return to play after anatomic anterior cruciate ligament reconstruction. *Clin Sports Med.* 2013;32(1):165-75. **PubMed | Google Scholar**
28. Feller J, Webster KE. Return to sport following anterior cruciate ligament reconstruction. *Int Orthop.* 2013;37(2):285-90. **PubMed | Google Scholar**
29. Tandogan RN, Taser O, Kayaalp A, Taskiran E, Pinar H, Alparslan B, Alturfan A. Analysis of meniscal and chondral lesions accompanying anterior cruciate ligament tears: relationship with age, time from injury, and level of sport. *Knee Surg Sports Traumatol Arthrosc.* 2004;12(4):262-70. **PubMed | Google Scholar**

30. Church S, Keating JF. Reconstruction of the anterior cruciate ligament: timing of surgery and the incidence of meniscal tears and degenerative change. *J Bone Joint Surg Br.* 2005;87(12):1639-42. **PubMed | Google Scholar**
31. Granan LP, Bahr R, Lie SA, Engebretsen L. Timing of anterior cruciate ligament reconstructive surgery and risk of cartilage lesions and meniscal tears: a cohort study based on the Norwegian National Knee Ligament Registry. *Am J Sports Med.* 2009;37(5):955-61. **PubMed | Google Scholar**
32. Demirag B, Aydemir F, Danis M, Ermutlu C. Incidence of meniscal and osteochondral lesions in patients undergoing delayed anterior cruciate ligament reconstruction. *Acta Orthop Traumatol Turc.* 2011;45(5):348-52. **PubMed | Google Scholar**
33. Gustavsson A, Neeter C, Thomee P, Silbernagel KG, Augustsson J, Thomee R, Karlsson J. A test battery for evaluating hop performance in patients with an ACL injury and patients who have undergone ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2006; 14(8):778-88. **PubMed | Google Scholar**
34. Myer GD, Paterno MV, Ford KR, Hewett TE. Neuromuscular training techniques to target deficits before return to sport after anterior cruciate ligament reconstruction. *J Strength Cond Res.* 2008;22(3):987-1014. **PubMed | Google Scholar**
35. Borsa PA, Lemhart SM, Irrgang JJ, Safran MR, Fu FH. The effects of joint position and direction of joint motion on proprioceptive sensibility in anterior cruciate ligament-deficient athletes. *Am j sports Med.* 1997;25(3):336-40. **PubMed | Google Scholar**

Table 1: Knee scores

Knee scores	Preoperative	Postoperative	P
Lysholm scores	60,7±12.5	90,3±4.8	<0.05
Tegner activity scores	3,52±1,4	6.2±1.5	<0.05
Cincinnati scores	14.8±5.3	26.9±1.6	<0.05

Table 2: Comparison of outcomes between work-accident and sports injury patients

Variables	Work-accident	Sports injury	P
N	16	17	
Preoperative Lysholm scores	59.1±11.2 (34-77)	62.1±13.8 (42-84)	0,417
Postoperative Lysholm scores	87.5±5.2 (78-95)	92.8±2.6 (90-100)	0,003
Preoperative Tegner activity scores	3.4±1.5 (1-6)	3.5±1.4 (2-7)	0,768
Postoperative Tegner activity scores	5.3±1.2 (3-7)	7.0±1.4 (5-9)	0,003
Preoperative Cincinnati scores	14.0±4.7 (8-24)	15.7±5.9 (6-26)	0,376
Postoperative Cincinnati scores	26.0±1.5 (24-29)	27.7±1.2 (26-30)	0,003
ROM deficits	-8.8±4.7 (-15-0)	-6.7±3.3 (-10-0)	0,001
Average time for returning to work	17.6±4.4 (12-27)	13.0±1.5 (10-15)	0,001

N=Number of patients, Postoperative and preoperative knee scores±standart deviation (Min-Max), ROM=Range of motion

Table 3: Comparison of outcomes between patients with or without meniscus lesions			
Variables	With meniscus lesion	No meniscus lesion	P
N	25	8	
Preoperative Lysholm scores	59.9±11.4 (34-80)	63.0±16.0 (43-84)	NS
Postoperative Lysholm scores	90.3±4.9 (78-100)	90.2±4.9 (80-95)	NS
Preoperative Tegner activity scores	3.4±1.4 (1-6)	3.6±1.5 (2-7)	NS
Postoperative Tegner activity scores	5.8±1.4(3-9)	7.3±1.5 (5-9)	NS
Preoperative Cincinnati scores	14.5±5.3 (6-26)	16.0±5.5 (8-24)	NS
Postoperative Cincinnati scores	26.9±1.7 (24-30)	27.0±0.9 (26-28)	NS
Average time for returning to work	15.3±4.3 (10-27)	15.2±2.9 (12-21)	NS
Postoperative and preoperative knee scores±SD (Min-Max). N=Number of patients, NS=Nonsignificant ($P>0.05$), SD=Standard deviation			