

Macrofungi of Türkmenbaba Mountain (Eskişehir)

Hasan KÖSTEKİ

Eskişehir Osmangazi University, The Graduate School of Natural and Applied Science, Eskişehir - TURKEY

Mustafa YAMAÇ

Eskişehir Osmangazi University, Science and Arts Faculty, Department of Biology, Eskişehir - TURKEY

M. Halil SOLAK

Muğla University, Ula Ali Koçman Vocational High School, Program of Fungi, Muğla - TURKEY

Received: 28.04.2004

Accepted: 08.09.2005

Abstract: The macrofungi collected from the Türkmenbaba mountain area in 2001-2003 were investigated. As a result of field and laboratory studies, 84 taxa belonging to 30 families were identified. Seven species belong to the class *Ascomycetes* and 77 species to *Basidiomycetes*. Seven of these taxa are new records for the Turkish macromycota.

Key Words: Macrofungi, Turkish Macromycota, Türkmenbaba Mountain

Türkmenbaba Dağı (Eskişehir) Makrofungalıları

Özet: Bu çalışmada, Türkmenbaba Dağında 2001-2003 yılları arasında yapılan arazi çalışmaları ile toplanan makrofungalılar araştırılmıştır. Arazi ve laboratuvar çalışmaları sonucunda 30 familyaya ait 84 takson belirlenmiştir. Belirlenen taksonlardan 7'si *Ascomycetes*, 77'si *Basidiomycetes* sınıfına aittir. Bu taksonlardan 7 tanesi Türkiye makromikotası için yeni kayittır.

Anahtar Sözcükler: Makrofungalılar, Türkiye Makromikotası, Türkmenbaba Dağı

Introduction

Although many studies have been carried out on the macrofungi flora of Turkey, there are numerous regions which have not been investigated yet. Türkmenbaba mountain, chosen as the study area, is near the Eskişehir-Kütahya highway, 25 km from Eskişehir (Figure 1). *Pinus nigra* J.F.Arnold, *Pinus sylvestris* L. and *Quercus cerris* L. are very common plants in the region. Climatic factors, plant distributions and field features of this area provide appropriate conditions for mushroom growth. Based on a review of the relevant literature, it appears that no previous macrofungi studies have been carried out in this area. The aims of this study were to determine the macrofungi taxa of this district, to identify edible, inedible and poisonous species and to make a contribution to the macrofungal flora of Turkey.

Materials and Methods

During field trips to Türkmenbaba mountain, macrofungi samples were collected periodically during all

seasons between 2001 and 2003. All species were photographed and their morphological and ecological features and localities were noted in the field. Specimens were then taken to the laboratory. Spore prints were obtained and spore dimensions were measured using an ocular micrometer. Samples were dried under suitable conditions and preserved in polyethylene bags containing thymol crystal. Identification of the specimens was carried out according to Phillips (1981), Moser (1983), Breitenbach and Kränzlin (1984, 1986, 1991, 1995), Cetto (1988) and Ellis and Ellis (1990). After identification of the specimens, some comparative studies were carried out on the macrofungi flora of Turkey (Baytop, 1994; Mat & Güçin, 2000; Afyon & Konuk, 2001; Doğan et al., 2001; Huseyinov et al., 2001; Kaşık et al., 2001; Kaya, 2001; Öztürk et al., 2001; Solak et al., 2001; Afyon & Konuk, 2002; Asan et al., 2002; Demirel et al., 2002; Doğan et al., 2002; Kaşık et al., 2002; Öztürk, 2002; Öztürk et al., 2002; Solak & Yılmaz, 2002; Solak et al., 2002; Yılmaz & İşiloğlu, 2002; Aktaş et al., 2003; Demirel et al., 2003; Doğan et




Figure 1. Map of the study area.

al., 2003; Kaşik et al., 2003; Öztürk et al., 2003; Pekşen & Karaca, 2003; Solak & Yılmaz Ersel, 2003; Solak et al., 2003; Türkekul, 2003; Türkekul & Sesli, 2003; Yılmaz Ersel et al., 2003; Afyon & Yağız, 2004; Demirel & Uzun, 2004; Demirel et al., 2004; Kaya et al., 2004; Yılmaz Ersel & Solak, 2004; Uzun et al., 2004). The specimens were prepared as herbarium materials and stored in the Fungiculture laboratory at the Department of Biology, Eskişehir Osmangazi University.

Results

Species determined in the research area are listed below. The numbers refer to the herbarium numbers and

an asterisk (*) means that the taxon is new for the Turkish macromycota.

ASCOMYCETES

Helvellaceae

1. *Gyromitra esculenta* (Pers.) Fr.

Hazardere district, in pine forest, 1110 m, 31.05.2003, Köstekci 590.

2. *Gyromitra infula* (Schaeff.) Quél.

Efsunbaba district, in mixed woodlands, 1150 m, 23.11.2002, Köstekci 481.

- 3. Helvella acetabulum (L.) Quél.**
Syn: *Peziza acetabulum* L.
Paxina acetabulum (L.) Kuntze
Acetabula vulgaris Fuckel
Kaklık deresi district, in meadows, 1102 m, 28.04.2003, Köstekci 576.
- 4. Helvella lacunosa Afzel.**
Hazardere district, in meadows, 1170 m, 28.10.2002, Köstekci 315.
- Humariaceae**
- 5. Aleuria aurantia (Pers.) Fuckel**
Hazardere district, near stream, 1170 m, 19.10.2002, Köstekci 282.
- Pezizaceae**
- *6. Peziza violacea Pers.**
Hazardere district, after fire in burnt sites, 1170 m, 05.04.2003, Köstekci 548.
- Diatrypaceae**
- 7. Diatrype disciformis (Hoffm.) Fr.**
Efsunbaba district, on deciduous tree, 1150 m, 23.11.2002, Köstekci 489; Hazardere district, 1170 m, 04.12.2002, Köstekci 530.
- BASIDIOMYCETES**
- Tremellaceae**
- 8. Exidia truncata Fr.**
Karatepe district, on deciduous tree, 1348 m, 24.03.2002, Köstekci 08.
- Nidulariaceae**
- 9. Crucibulum laeve (Huds.) Kambly**
Syn: *Crucibulum vulgare* Tul. & C.Tul.
Kaklık deresi district, on deciduous tree, 1102 m, 28.09.2002, Köstekci 166.
- 10. Cyathus olla (Batsch) Pers.**
Kaklık deresi district, on deciduous tree, 1102 m, 28.09.2002, Köstekci 165.
- Sphaerobolaceae**
- *11. Sphaerobolus stellatus Tode**
Hazardere district, on deciduous tree, 1170 m, 19.10.2002, Köstekci 283.
- Rhizopogonaceae**
- 12. Rhizopogon luteolus Fr.**
Kaklık deresi district, in mixed woodlands, 1102 m, 12.10.2002, Köstekci 210.
- 13. Rhizopogon roseolus (Corda) Th.Fr.**
Karatepe district, in mixed woodlands, 1217 m, 21.04.2002, Köstekci 21.
- Lycoperdaceae**
- 14. Handkea excipuliformis (Scop.) Kreisel**
Syn: *Calvatia excipuliformis* (Scop.) Perdeck
Calvatia saccata (Vahl) Morgan
Hazardere district, in meadows, 1170 m, 31.05.2003, Köstekci 597.
- 15. Lycoperdon pyriforme Schaeff.**
Kaklık deresi district, on pine stumps, 1102 m, 28.09.2002, Köstekci 156.
- 16. Lycoperdon perlatum Pers.**
Syn: *Lycoperdon gemmatum* Batsch
Kaklık deresi district, in meadows, 1102 m, 28.09.2002, Köstekci 159.
- *17. Lycoperdon pedicellatum Peck**
Kaklık deresi district, in meadows, 1102 m, 28.09.2002, Köstekci 502.
- 18. Lycoperdon umbrinum Pers.**
Kaklık deresi district, in meadows, 1102 m, 28.09.2002, Köstekci 130.
- 19. Bovista plumbea Pers.**
Kaklık deresi district, in meadows, 1102 m, 28.09.2002, Köstekci 136; Kaklık deresi district, 1130 m, 12.10.2002, Köstekci 224; Yukarı Kalabak village district, 1160 m, 16.11.2002, Köstekci 430.
- *20. Bovista aestivalis (Bonord.) Demoulin**
Keltepe district, in meadows, 1110 m, 02.11.2002, Köstekci 395.
- Sclerodermataceae**
- 21. Scleroderma cepa Pers.**
Efsunbaba district, near stream, 1560 m, 07.07.2002, Köstekci 96.

22. *Scleroderma verrucosum* (Bull.) Pers.
Kaklık deresi district, under pine trees, 1102 m, 28.04.2003, Köstekci 579.
23. *Scleroderma areolatum* Ehrenb.
Kaklık deresi district, under deciduous tree, 1102 m, 12.10.2002, Köstekci 207.
- Thelephoraceae
24. *Boletopsis leucomelaena* (Pers.) Fayod
Hızadere district, in pine forest, 1170 m, 19.10.2002, Köstekci 271.
25. *Sarcodon imbricatus* (L.) P.Karst.
Kaklık deresi district, near stream, 1130 m, 28.09.2002, Köstekci 117, 1102 m, 12.10.2002, Köstekci 261.
- Boletaceae
26. *Boletus edulis* Bull.
Kaklık deresi district, in mixed woodlands, 1102 m, 12.10.2002, Köstekci 236.
27. *Boletus chrysenteron* Bull.
Syn: *Xerocomus chrysenteron* (Bull.) Quél.
Kaklık deresi district, in mixed woodlands, 1110 m, 28.09.2002, Köstekci 185.
28. *Suillus luteus* (L.) Gray
Kaklık deresi district, in pine forest, 1102 m, 12.10.2002, Köstekci 210.
29. *Suillus bovinus* (Pers.) Kuntze
Syn: *Ixocomus bovinus* (L.) Quél.
Hızadere district, in pine forest, 1170 m, 28.10.2002, Köstekci 309.
30. *Suillus bellinii* (Inzenga) Watling
Kaklık deresi district, in pine forest, 1102 m, 12.10.2002, Köstekci 231.
- Hymenochaetaceae
31. *Phellinus pini* (Brot.) Bondartsev & Singer
Kaklık deresi district, on wood of live pine, 1102 m, 28.09.2002, Köstekci 186.
- Polyporaceae
- *32. *Albatrellus subrubescens* (Murrill) Pouzar
Hızadere district, in pine forest, 1170 m, 28.10.2002, Köstekci 320.
33. *Pycnoporus cinnabarinus* (Jacq.) Fr.
Syn: *Trametes cinnabarina* (Jacq.) Fr.
Kaklık deresi district, on deciduous tree, 1102 m, 28.09.2002, Köstekci 184.
34. *Gloeophyllum abietinum* (Bull.) P.Karst.
Syn: *Lenzites abietina* (Bull.) Fr.
Kaklık deresi district, on pine stumps, 1102 m, 19.04.2003, Köstekci 564.
35. *Polyporus varius* (Pers.) Fr.
Efsunbaba district, on pine, 1150 m, 23.11.2002, Köstekci 474.
36. *Polyporus arcularius* (Batsch) Fr.
Syn: *Polyporus anisoporus* Delastre & Mont.
Efsunbaba district, on broadleaved tree, 1230 m, 23.11.2002, Köstekci 438.
37. *Trametes versicolor* (L.) Lloyd
Syn: *Coriolus versicolor* (L.) Quél.
Kaklık deresi district, on poplar stumps, 1102 m, 28.09.2002, Köstekci 188.
38. *Trametes hirsuta* (Wulfen) Pilát
Kaklık deresi district, on deciduous tree, 1102 m, 28.04.2003, Köstekci 587.
- Hydnaceae
39. *Hydnum repandum* L.
Syn: *Dentinum repandum* (L.) Gray
Kaklık deresi district, in pine forest, 1102 m, 26.10.2002, Köstekci 295.
- Corticaceae
40. *Stereum hirsutum* (Willd.) Pers.
Domuzgölü hill, on pine stumps, 1250 m, 16.03.2002, Köstekci 07.
- Ramariaceae
41. *Ramaria fumigata* (Peck) Corner
Syn: *Ramaria versatilis* Quél.
Hızadere district, in pine forest, 1170 m, 28.10.2002, Köstekci 321.
42. *Ramaria flava* (Tourn. ex Battarra) Quél.
Syn: *Clavaria flava* (Tourn. ex Battarra) Schaeff.

Kaklık deresi district, in pine forest, 1102 m, 12.10.2002, Köstekci 211.

Clavariaceae

43. Clavariadelphus truncatus (Quél.) Donk

Kirazlı road district, in mixed woodlands, 1102 m, 13.10.2002, Köstekci 192.

44. Clavariadelphus pistillaris (L.) Donk

Efsunbaba district, in mixed woodlands, 1560 m, 23.11.2002, Köstekci 495.

Schizophyllaceae

45. Schizophyllum commune Fr.

Keltepe district, on pine stumps, 1110 m, 02.11.2002, Köstekci 365.

Entolomataceae

46. Entoloma hirtipes (Schumach.) M.M.Moser

Kaklık deresi district, in pine forest, 1102 m, 28.04.2003, Köstekci 568.

Russulaceae

47. Russula delica Fr.

Kaklık deresi district, in pine forest, 1102 m, 12.10.2002, Köstekci 194.

48. Lactarius chrysorrheus Fr.

Efsunbaba district, under Quercus trees, 1560 m, 23.11.2002, Köstekci 482.

49. Lactarius deliciosus (L.) Gray

Kaklık deresi district, in pine forest, 1102 m, 28.09.2002, Köstekci 133; 12.10.2002, Köstekci 193.

50. Lactarius deterrimus Gröger

Kaklık deresi district, in pine forest, 1102 m, 12.10.2002, Köstekci 191.

Coprinaceae

51. Coprinus comatus (O.F.Müll.) Gray

Kaklık deresi district, in meadows, near cow dung, 1102 m, 12.10.2002, Köstekci 237.

52. Coprinus micaceus (Bull.) Fr.

Keltepe district, on rotten poplar stumps, 1115 m, 31.05.2003, Köstekci 620.

Gomphidiaceae

53. Chroogomphus rutilus (Schaeff.) O.K.Mill.

Syn: *Gomphidius viscidus* (L.) Fr.

Gomphidius rutilus (Schaeff.) S.Lundell

Kaklık deresi district, in pine forest, 1102 m, 28.09.2002, Köstekci 116-131.

Strophariaceae

54. Hypholoma fasciculare (Huds.) Quél.

Syn: *Nematoloma fasciculare* (Huds.) P.Karst.

Kaklık deresi district, on pine stumps, 1120 m, 28.09.2002, Köstekci 162.

55. Stropharia aeruginosa (Curtis) Quél.

Hazardere district, in mixed woodlands, 1170 m, 28.10.2002, Köstekci 313.

56. Stropharia caerulea Kreisel

Keltepe district, in meadows, 1110 m, 02.11.2002, Köstekci 345.

Agaricaceae

57. Agaricus impudicus (Rea) Pilát

Kaklık deresi district, near road, in meadows, 1102 m, 28.09.2002, Köstekci 148.

Paxillaceae

58. Paxillus involutus (Batsch) Fr.

Kirazlı road district, the bank of stream, 1210 m, 28.09.2002, Köstekci 122; Kaklık deresi district, 1102 m, 12.10.2002, Köstekci 146; Keltepe district, 1250 m, 26.10.2002, Köstekci 246.

Hygrophoraceae

59. Hygrocybe nigrescens (Quél.) Kühner

Kaklık deresi district, in meadows, 1102 m, 12.10.2002, Köstekci 205.

60. Hygrophorus chrysodon (Batsch) Fr.

Hazardere district, in pine forest, 1170 m, 19.10.2002, Köstekci 296; Yukarı Kalabak village, 1260 m, 16.11.2002, Köstekci 430, Karatepe district, 1230 m, 30.11.2002, Köstekci 503.

61. Hygrophorus agathosmus (Fr.) Fr.

Keltepe district, under deciduous tree, 1250 m, 02.11.2002, Köstekci 267.

62. Hygrophorus hypothejus (Fr.) Fr.

Hazardere district, in mixed woodlands, 1170 m, 28.10.2002, Köstekci 319.

63. *Hygrophorus camarophyllus* (Alb. & Schwein.)
Dumée, Grandjean & Maire

Syn: *Hygrophorus caprinus* (Scop.) Fr.

Hazardere district, in pine forest, 1170 m,
19.10.2002, Köstekci 265.

64. *Hygrophorus discoideus* (Pers.) Fr.

Kaklık deresi district, in pine forest, 1102 m,
12.10.2002, Köstekci 204.

Amanitaceae

65. *Amanita caesarea* (Scop.) Pers.

Kaklık deresi district, in pine forest, 1102 m,
28.09.2002, Köstekci 118.

66. *Amanita pantherina* (DC.: Fr.) Krombh.

Kaklık deresi district, in pine forest, 1102 m,
28.09.2002, Köstekci 120. Kaklık deresi district, 1102
m, 12.10.2002, Köstekci 195.

Lepiotaceae

67. *Leucoagaricus leucothites* (Vittad.) M.M.Moser
ex Bon

Syn: *Leucoagaricus pudicus* (Bull.) Bon

Kaklık deresi district, in meadows, 1102 m,
28.09.2002, Köstekci 121.

68. *Macrolepiota procera* (Scop.) Singer

Kaklık deresi district, in pine forest, 1102 m,
28.09.2002, Köstekci 183; Kaklık deresi district, 1120
m, 19.04.2003, Köstekci 591.

69. *Macrolepiota excoriata* (Schaeff.) M.M.Moser

Keltepe district, in pine forest, 1110 m, 02.11.2002,
Köstekci 329.

Tricholomataceae

*70. *Strobilurus tenacellus* (Pers.) Singer

Syn: *Pseudohiatula tenacella* (Pers.) Métrod

Collybia tenacella (Pers.) P.Kumm.

Kaklık deresi district, in pine forest, 1102 m,
21.04.2003, Köstekci 572.

71. *Melanoleuca graminicola* (Velen.) Kühner &
Maire

Syn: *Tricholoma graminicola* Velen.

Hazardere district, in meadows, 1170 m,
19.10.2002, Köstekci 303.

72. *Pseudoclitocybe cyathiformis* (Bull.) Singer

Kaklık deresi district, in pine forest, 1102 m,
12.10.2002, Köstekci 248; Efsunbaba district, 1560 m,
23.11.2002, Köstekci 462.

73. *Lepista nuda* (Bull.) Cooke

Syn: *Tricholoma nudum* (Bull.) P.Kumm.

Rhodopaxillus nudus (Bull.) Maire.

Keltepe district, in pine forest, 1230 m, 02.11.2002,
Köstekci 373.

74. *Cystoderma carcharias* (Pers.) Fayod

Keltepe district, in pine forest, 1110 m, 02.11.2002,
Köstekci 369.

75. *Cystoderma amianthinum* (Scop.) Fayod

Hazardere district, in pine forest, 1170 m,
19.10.2002, Köstekci 307.

76. *Clitocybe geotropa* (Bull.) Quél.

Kirazlı road district, in pine forest, 1210 m,
13.10.2002, Köstekci 203.

77. *Mycena epipterygia* (Scop.) Gray

Hazardere district, in pine forest, 1170 m,
19.10.2002, Köstekci 258.

78. *Mycena polygramma* (Bull.) Gray

Keltepe district, under deciduous trees, 1110 m,
02.11.2002, Köstekci 331.

79. *Marasmius androsaceus* (L.) Fr.

Kaklık deresi district, on pine twigs, 1102 m,
12.10.2002, Köstekci 228, Keltepe district, 1110 m,
02.11.2002, Köstekci 367, Yukarı Kalabak village
district, 1250 m, 16.11.2002, Köstekci 407.

*80. *Marasmius recubans* Quél.

Keltepe district, on Quercus sp. leaves, 1110 m,
02.11.2002, Köstekci 366.

81. *Tricholoma equestre* (L.) P.Kumm.

Syn: *Tricholoma auratum* (Paulet) Gillet

Kaklık deresi district, in pine forest, 1125 m,
28.09.2002, Köstekci 174.

82. *Tricholoma fracticum* (Britzelm.) Kreisel

Syn: *Tricholoma batschii* Gulden

Tricholoma subannulatum (Batsch) Bres.

Keltepe district, in pine forest, 1240 m, 02.11.2002, Köstekci 255.

83. Collybia dryophila (Bull.) P.Kumm.

Karatepe district, in mixed woodlands, 1150 m, 30.11.2002, Köstekci 498.

84. Armillariella mellea (Vahl) P.Kumm.

Kirazlı road district, in mixed woodlands, 1160 m, 12.10.2002, Köstekci 227.

Discussion and Conclusion

In this study, 84 species of macrofungi were identified, 7 of which belong to the class *Ascomycetes* and 77 to *Basidiomycetes*. When our results are compared with other studies carried out near the research area, the similarity between the mycota of Türkmenbaba mountain and neighbouring regions is not significant. Only 10 species determined in the present study were previously

reported from Ankara (Watling & Gregory, 1977; Özdal & İlbay, 2000; Güler & Mutlu, 2002) and 12 species from Bolu (Lohwag & Ünlügil, 1957; Niemela & Uotila, 1977; Watling & Gregory, 1977; Sümer, 1982). This could be because of differences in vegetation, climate and flora. Forty-six of the 84 macrofungus species found in the area are considered edible, 9 are poisonous and 29 are inedible.

With this study, 7 species (*Peziza violacea*, *Sphaerobolus stellatus*, *Lycoperdon pedicellatum*, *Bovista aestivalis*, *Albatrellus subrubescens*, *Strobilurus tenacellus* and *Marasmius recubans*) were added to the Turkish macromycota as new records.

Acknowledgements

The authors would like to thank the Scientific Research and Application Centre (FBAM) of Osmangazi University, Eskişehir, for its financial support.

References

- Afyon A & Konuk M (2001). Batı Karadeniz Bölgesinde halkın tanıldığı bazı önemli yenen mantarlar. *Selçuk Univ Fen Bilimleri Dergisi* 9: 109-118.
- Afyon A & Konuk M (2002). Zonguldak Yöresi makrofungalıları üzerine bir araştırma. *Ot Sistematisk Botanik Dergisi* 9(1): 121-128.
- Afyon A & Yağız D (2004). Macrofungi of Sinop province Turkey. *Turk J Bot* 28: 351-360.
- Aktaş S, Öztürk C, Kaşık G, Sabahlar Ş & Doğan HH (2003). Macrofungi flora of Bozkır district (Konya). *Turk J Bot* 27: 37-43.
- Asan A, Sesli E, Güçin F & Stoichev G (2002). *Myriostoma coliforme* and *Phyllophoria ribis* (*Basidiomycetes*): First reports from European Turkey. *Botanika Chronika* 15: 45-49.
- Baytop A (1994). Türkiye' nin makrofungalıları ile ilgili bir yayın listesi. *Turk J Bot* 18: 175-185.
- Breitenbach J & Kränzlin F (1984). *Fungi of Switzerland, Volume 1. Ascomycetes*, Luzern: Verlag mycologia.
- Breitenbach J & Kränzlin F (1986). *Fungi of Switzerland, Volume 2. Nongilled Fungi*, Luzern: Verlag mycologia.
- Breitenbach J & Kränzlin F (1991). *Fungi of Switzerland, Volume 3. Boletes and Agarics 1st part*, Luzern: Verlag mycologia.
- Breitenbach J & Kränzlin F (1995). *Fungi of Switzerland, Volume 4. Agarics 2nd part*, Luzern: Verlag mycologia.
- Cetto B (1988). *Enzyklopädie der Pilze*, Band 3., München: BLV Verlagsgesellschaft
- Demirel K, Kaya A & Uzun Y (2003). Macrofungi of Erzurum province. *Turk J Bot* 27: 29-36.
- Demirel K & Uzun Y (2004). Two new records of Phallales for the mycoflora of Turkey. *Turk J Bot* 28: 213 – 214.
- Demirel K, Uzun Y & Kaya A (2002). Macrofungi of Ağrı province. *Turk J Bot* 26: 291-295.
- Demirel K, Uzun Y & Kaya A (2004). Some poisonous fungi of East Anatolia. *Turk J Bot* 28: 215-219.
- Doğan HH, Gürer M & Öztürk C (2001). Two new Ascomytes genus for the fungal flora of Turkey. *Ot Sistematisk Botanik Dergisi* 8(1): 113-118.
- Doğan HH, Öztürk C & Kaşık G (2002). Two new *Peziza* records from Mut environ. *Ot Sistematisk Botanik Dergisi* 9(2): 117-122.
- Doğan HH, Öztürk C, Kaşık G, & Aktaş S (2003). New records for the mycoflora of Turkey from Mut environ. *Ot Sistematisk Botanik Dergisi* 10(2): 197-211.
- Ellis MB & Ellis CP (1990). *Fungi without gills (Hymenomycetes and Gasteromycetes)*, London: Chapman and Hill.
- Huseyinov E, Selçuk F & Aslantaş I (2001). Some data on Agaricoid fungi from Sivas province (Turkey). *Mycology & Phytopathology* 35: 29-33.
- Kaşık G, Doğan HH, Öztürk C & Aktaş S (2003). New records of Tricholomataceae and Cortinariaceae for Turkish macrofungi flora from Alanya (Antalya) District. *Ot Sistematisk Botanik Dergisi* 10(1): 143-168.
- Kaşık G, Öztürk C & Toprak E (2001). Macrofungi of Niğde province (Turkey). *Ot Sistematisk Botanik Dergisi* 8(2): 137-142.

- Kaşık G, Öztürk C, Türkoğlu A & Doğan HH (2002). Macrofungi flora of Yeşilhisar district (Kayseri). *Ot Sistematisk Botanik Dergisi* 9(2): 123-134.
- Kaya A (2001). Contributions to the macrofungi flora of Bitlis province, *Tr J of Botany* 25: 379-383.
- Kaya A, Akan Z, Demirel K (2004). A checklist of macrofungi of Besni (Adiyaman) District, *Turk J Bot* 28: 247-251.
- Lohwag K & Ünlügil H (1957). Türkiye'nin mantar florası hakkında araştırma. *İ.Ü.Orman Fakültesi Orman Botanığı Enstitüsü Dergisi* 129:137.
- Mat A & Gücin F (2000). Türkiye' nin makrofungalıları ile ilgili bir yayın listesi (Ek liste 2), In: Mat A (ed.) *Türkiye'de Mantar Zehirlenmeleri ve Zehirli Mantarlar*, pp. 212-217, Ankara: Nobel.
- Moser M (1983). *Keys to Agarics and Boleti*, Stuttgart: Gustav, Fischer Verlag
- Niemela T & Uotila P (1977). Lignicolous macrofungi from Turkey and Iran. *Karstenia* 17: 33-39.
- Özdal S & İlbay E (2000). Ankara sınırları içinde seçilmiş orman alanlarında doğal olarak yetişen şapkalı mantarların tanınması, In: Eltez S & Eren E (eds.) *VI. Yemeklik Mantar Kongresi*, pp. 168-185, İstanbul: Ömür matbaacılık.
- ÖzTÜRK C (2002). Türkiye makrofungus florası için iki yeni kayıt. *Ot Sistematisk Botanik Dergisi* 9(1): 117-120.
- ÖzTÜRK C, Gürer M & Doğan HH (2001). Two new records for the fungal flora of Turkey. *Ot Sistematisk Botanik Dergisi* 8(2): 133-136.
- ÖzTÜRK C, Doğan HH, Aktaş S & Kaşık G (2002). New records for the macrofungi Flora of Turkey from Ahırlı and Yalıhüyük districts (Konya). *Ot Sistematisk Botanik Dergisi* 9(2): 135-148.
- ÖzTÜRK C, Doğan HH, Kaşık G & Aktaş S (2003). Türkiye mikoforası için Karaman yöresinde yeni kayıtlar. *Ot Sistematisk Botanik Dergisi* 10(2): 213-248.
- Pekşen A & Karaca G (2003). Macrofungi of Samsun province. *Turk J Bot* 27: 173-184.
- Phillips R (1981). *Mushroom and other fungi of Great Britain and Europe*, London: Pan Books Ltd.
- Solak MH & Yılmaz F (2002). Manisa yöresi makrofungus florasına katkılar. *Ekoloji Çevre Dergisi* 10(43): 30-32.
- Solak MH & Yılmaz Ersel F (2003). Muğla yöresinden Türkiye makromikotasına yeni kayıtlar. *Ekoloji Çevre Dergisi* 10(48): 10-12.
- Solak MH, Gücin F, Yılmaz F & İşiloğlu M (2003). Some macrofungi from Çanakkale province. *Ot Sistematisk Botanik Dergisi* 10(1): 97-109.
- Solak MH, Kalılmış E & İşiloğlu M (2001). New records for the fungi flora of Turkey. *Bio-Science Research Bulletin* 17: 99-103.
- Solak MH, Yılmaz Ersel F, Gücin F & İşiloğlu M (2002). Macrofungi of Balıkesir province from Turkey, *Bio-Science Research Bulletin* 18(2): 137-149.
- Sümer S (1982). *Batı Karadeniz bölgesi özellikle Bolu çevresinde bulunan odun tahirçisi mantarlar*. İstanbul: İ.Ü. Orman Fak. Yayınları.
- Türkekul İ (2003). A contribution to the fungal flora of Tokat province. *Turk J Bot* 27: 313-320.
- Türkekul İ & Sesli E (2003). Macrofungi of Gümenek picnic area of Tokat province. *Bio-Science Research Bulletin* 19(2): 117-120.
- Uzun Y, Keleş A, Demirel K & Solak MH (2004). Some macrofungi from Bayburt province in Turkey. *Bulletin of Pure and Applied Sciences* 23B(1): 47-55.
- Watling R & Gregory NM (1977). Larger fungi from Turkey, Iran and neighbouring countries. *Karstenia* 17: 59-72.
- Yılmaz F & İşiloğlu M (2002). Macrofungi of Değirmenboğazı (Balıkesir). *Turk J Bot* 26: 161-164.
- Yılmaz Ersel F & Solak MH (2004). Three new records for the Turkish macromycota. *Ekoloji* 13(52): 7-8.
- Yılmaz Ersel F, Solak MH & Gücin F (2003). Two new records from Anatolia. *Bio-Science Research Bulletin* 19(2): 83-85.