

On the Relations Between Fuzzy Topologies and α Cut Topologies

Zekeriya GÜNEY¹

Abstract: In this study, some relations have been generated between fuzzy sets and their cross-sections, and further relations have been derived between fuzzy topological spaces and α -cut topologies by using these relations.

Key words and phrases: Fuzzy set, α -cut, fuzzy topological spaces, α -cut topologies

Fuzzy Topolojiler ve α -kesit Topolojiler Arasındaki Bağlılıklar Üzerine

Özet: Bu çalışmada fuzzy kümeler ve bunların α -kesitleri arasında bazı bağıntılar üretilmiş ve bu bağıntıları kullanarak, fuzzy topolojik uzaylar ve α -kesit topolojileri arasında bazı ilişkiler ortaya koyulmuştur.

Anahtar sözcük ve deyimler: Fuzzy küme, α -kesit, fuzzy topolojik uzaylar, α -kesit topolojiler

Introduction

Standard symbols and deduction formulas of logistics have been used in the definitions, assumptions, and proofs that have been provided in this paper. Universal symbolic language of mathematics have been used instead of a national language, where applicable.

Let (X, \mathcal{F}) be a fuzzy topological space (f.t.s), A be a fuzzy set in space (X, \mathcal{F}) , $\alpha \in (0, 1] = I_0$, $A_\alpha = \{x \mid A(x) \geq \alpha\}$ be a α -cut of set A and $\mathcal{F}_\alpha = \{A_\alpha \mid A \in \mathcal{F}\}$ be the family of α -cuts of the elements of \mathcal{F} . It has been proven that family \mathcal{F}_α a basis for at least one topology on X , and some fuzzy-crisp relations have been derived by determining some sufficient conditions for X to be a topological space.

Preliminaries

($I := [0, 1]$) ($Y^X = \{f \mid f: X \rightarrow Y \text{ function}\}$)
($f \in Y^X$) ($A \in I^X$) ($B \in I^Y$) ($\mathcal{A} \subset I^X$) \Rightarrow

1. $(f^{-1}[B])(x) = B(f(x))$ [1]
2. $(\bigvee \mathcal{A})(x) := \sup \{A(x) \mid A \in \mathcal{A}\}$, $(\bigwedge \mathcal{A})(x) := \inf \{A(x) \mid A \in \mathcal{A}\}$,
3. (X, \mathcal{F}) f.t.s. \Leftrightarrow

($0 \in \mathcal{F}$) ($1 \in \mathcal{F}$) ($A, B \in \mathcal{F} \Rightarrow A \wedge B \in \mathcal{F}$) ($\mathcal{A} \subset \mathcal{F} \Rightarrow \bigvee \mathcal{A} \in \mathcal{F}$)
 A open $\Leftrightarrow A \in \mathcal{F}$, A closed $\Leftrightarrow \forall A \in \mathcal{F}$

¹ Muğla Üniversitesi Fen-Edebiyat Fakültesi Matematik Bölümü 48000- Muğla

4. $A^\circ = \vee \{ B \mid (B \leq A) (B \in \mathcal{F}) \}, \quad A \in \mathcal{F} \Leftrightarrow A = A^\circ$ [2]

5. \mathcal{B} base for $\mathcal{F} \quad \Leftrightarrow (\mathcal{B} \subset \mathcal{F}) [A \in \mathcal{F} \Rightarrow (\exists \mathcal{B}^* \subset \mathcal{B}) (A = \vee \mathcal{B}^*)]$ [3]

6. $(\alpha \in I) (A \in I^X) \Rightarrow A_\alpha = \{x \mid A(x) \geq \alpha\}, \quad \mathbf{0}_\alpha = \begin{cases} \emptyset, & \alpha \in (0,1) \\ X, & \alpha = 0 \end{cases}$

7. $\mathcal{A}_\alpha := \{A_\alpha \mid A \in \mathcal{A}\}, \quad |\mathcal{A}_\alpha| \leq |\mathcal{A}|$
 $f : (X, \mathcal{F}) \rightarrow (Y, \mathcal{E}) \Rightarrow$

8. f fuzzy continuous (f -c.) $\Leftrightarrow (A \in \mathcal{E} \Rightarrow f^{-1}[A] \in \mathcal{F})$

9. $(X, \mathcal{F}) T_0 \quad \Leftrightarrow$

$(\forall x, y \in X) (\forall \alpha \in I_0) (\exists A \in \mathcal{F}) [(P_x^\alpha \in A) (P_y^\alpha \notin A) \vee (P_x^\alpha \notin A) (P_y^\alpha \in B)]$

10. $P \in I^X, P(x) = \begin{cases} \alpha, & x = a \\ 0, & x \neq a \end{cases} \quad \Leftrightarrow P = P_a^\alpha$
 $A \in I^X \quad \Rightarrow (P_x^\alpha \in A \Leftrightarrow \alpha \leq A(x))$

Some Results For α -cuts of Fuzzy Sets

$A, B \in I^X, \alpha \in I, \mathcal{A} \subset I^X \Rightarrow$

- 1. $A \leq B \Leftrightarrow A_\alpha \subset B_\alpha, \quad A = B \Leftrightarrow A_\alpha = B_\alpha$
- 2. $(A \vee B)_\alpha = A_\alpha \cup B_\alpha, \quad (A \wedge B)_\alpha = A_\alpha \cap B_\alpha$ [4]
- 3. $\cup \mathcal{A}_\alpha \subset (\vee \mathcal{A})_\alpha, \quad (\wedge \mathcal{A})_\alpha \subset \cap \mathcal{A}_\alpha$

Proof:

$A \in \mathcal{A} \Rightarrow (A \leq \vee \mathcal{A}) (\wedge \mathcal{A} \leq A)$
 $\Rightarrow (A_\alpha \subset (\vee \mathcal{A})_\alpha) ((\wedge \mathcal{A})_\alpha \subset A_\alpha) \Rightarrow \cup \mathcal{A}_\alpha \subset (\vee \mathcal{A})_\alpha,$
 $\Rightarrow (\cup \mathcal{A}_\alpha \subset (\vee \mathcal{A})_\alpha) ((\wedge \mathcal{A})_\alpha \subset \cap \mathcal{A}_\alpha)$

4. $(\exists \mathcal{A} \subset I^X) (\exists \alpha \in I) ((\vee \mathcal{A})_\alpha \neq \cup \mathcal{A}_\alpha),$
 $\exists \mathcal{A} \subset I^X) (\exists \alpha \in I) ((\wedge \mathcal{A})_\alpha \neq \cap \mathcal{A}_\alpha)$

Proof : (Example)

$X = \mathbb{R}, \quad a \in I \Rightarrow A^a = \{ (x, a) \mid x \in \mathbb{R} \}, \quad \mathcal{A} = \{ A^a \mid a \in (1/2, 2/3) \} \subset 2^X,$
 for $\alpha = 2/3 \quad \mathcal{A}_{2/3} = \{ (A^a)_{2/3} \mid a \in (1/2, 2/3) \} = \{ \emptyset \}, \quad \cup \mathcal{A}_{2/3} = \emptyset,$
 $\vee \mathcal{A} = A^{2/3} = \{ (x, 2/3) \mid x \in \mathbb{R} \}, \quad (\vee \mathcal{A})_{2/3} = \mathbb{R}, \dots$

5. $|\mathcal{A}| < \aleph_0 \Rightarrow ((\vee \mathcal{A})_\alpha = \cup \mathcal{A}_\alpha) ((\wedge \mathcal{A})_\alpha = \cap \mathcal{A}_\alpha)$

Proof

$x \in (\vee \mathcal{A})_\alpha \Leftrightarrow (\vee \mathcal{A})(x) \geq \alpha \Leftrightarrow \sup \{ A(x) \mid A \in \mathcal{A} \} \geq \alpha$
 $\stackrel{\text{hip.}}{\Leftrightarrow} \max \{ A(x) \mid A \in \mathcal{A} \} \geq \alpha \Leftrightarrow (\exists A \in \mathcal{A}) (A(x) \geq \alpha)$
 $\Leftrightarrow (\exists A \in \mathcal{A}) (x \in A_\alpha) \Leftrightarrow x \in \cup \mathcal{A}_\alpha,$
 $(\wedge \mathcal{A})_\alpha \stackrel{2.6}{=} \{ x \mid (\wedge \mathcal{A})(x) \geq \alpha \} \stackrel{2.2}{=} \{ x \mid \inf \{ A(x) \mid A \in \mathcal{A} \} \geq \alpha \}$
 $\stackrel{\text{hip.}}{=} \{ x \mid \min \{ A(x) \mid A \in \mathcal{A} \} \geq \alpha \} = \{ x \mid A \in \mathcal{A} \Rightarrow A(x) \geq \alpha \}$
 2.6.

$$= \{x \mid A \in \mathcal{A} \Rightarrow x \in A_\alpha\} = \bigcap \mathcal{A}_\alpha.$$

6. $\forall \mathcal{A} \in \mathcal{A} \Rightarrow (\forall \mathcal{A})_\alpha = \bigcup \mathcal{A}_\alpha$, $\wedge \mathcal{A} \in \mathcal{A} \Rightarrow (\wedge \mathcal{A})_\alpha = \bigcap \mathcal{A}_\alpha$

Proof :

$$\begin{aligned} \forall \mathcal{A} \in \mathcal{A} &\Rightarrow (\forall \mathcal{A})_\alpha \in \mathcal{A}_\alpha \Rightarrow (\forall \mathcal{A})_\alpha \subset \bigcup \mathcal{A}_\alpha \stackrel{3.3.}{\Rightarrow} (\forall \mathcal{A})_\alpha = \bigcup \mathcal{A}_\alpha \\ : & \\ \wedge \mathcal{A} \in \mathcal{A} &\Rightarrow (\wedge \mathcal{A})_\alpha \in \mathcal{A}_\alpha \Rightarrow (\wedge \mathcal{A})_\alpha \supset \bigcap \mathcal{A}_\alpha \stackrel{3.3.}{\Rightarrow} (\wedge \mathcal{A})_\alpha = \bigcap \mathcal{A}_\alpha \end{aligned}$$

7. $(\setminus A)_\alpha = \setminus A_{1-\alpha} \cup A^{-1}(1-\alpha)$

Proof :

$$\begin{aligned} x \in (\setminus A)_\alpha &\Leftrightarrow (\setminus A)(x) \geq \alpha \Leftrightarrow 1 - A(x) \geq \alpha \Leftrightarrow A(x) \leq 1 - \alpha \\ &\Leftrightarrow A(x) < 1 - \alpha \vee A(x) = 1 - \alpha \Leftrightarrow x \notin A_{1-\alpha} \vee x \in A^{-1}(1-\alpha) \\ &\Leftrightarrow x \in \setminus A_{1-\alpha} \vee x \in A^{-1}(1-\alpha) \Leftrightarrow x \in \setminus A_{1-\alpha} \cup A^{-1}(1-\alpha). \end{aligned}$$

8. $(A \setminus B)_\alpha = (A_\alpha \setminus B_{1-\alpha}) \cup (A_\alpha \cap B^{-1}(1-\alpha))$

[4]

9. $(f \in Y^X)(A \in I^Y)(\alpha \in I) \Rightarrow (f^{-1}[A])_\alpha = f^{-1}[A_\alpha]$

Proof :

$$\begin{aligned} x \in (f^{-1}[A])_\alpha &\stackrel{2.6}{\Leftrightarrow} (f^{-1}[A])(x) \geq \alpha \stackrel{2.1.}{\Leftrightarrow} A(f(x)) \geq \alpha \\ &\stackrel{2.6.}{\Leftrightarrow} f(x) \in A_\alpha \Leftrightarrow x \in f^{-1}[A_\alpha] \end{aligned}$$

Some Relations Between Fuzzy Topologies and α -Cut Topologies:

1. (X, \mathcal{F}) f.t.s. $\Rightarrow (\forall \alpha \in I_0)(\exists (X, \mathcal{E})$ t.s.) $(\mathcal{F}_\alpha$ base for $\mathcal{E})$

Proof :

$$\begin{aligned} (X, \mathcal{F}) \text{ f.t.s.} &\stackrel{2.3}{\Rightarrow} 1 \in \mathcal{F} \\ &\stackrel{2.6-2.7}{\Rightarrow} (\forall \alpha \in I_0)(1_\alpha = X \in \mathcal{F}_\alpha) \dots (a) \\ A, B \in \mathcal{F}_\alpha &\stackrel{2.7.}{\Rightarrow} (\forall \alpha \in I_0)(\exists C, D \in \mathcal{F})(A = C_\alpha)(B = D_\alpha) \\ &\stackrel{\text{hip.} - 2.3.}{\Rightarrow} C \wedge D \in \mathcal{F} \stackrel{2.7.}{\Rightarrow} (C \wedge D)_\alpha \in \mathcal{F}_\alpha \\ &\stackrel{3.2.}{\Rightarrow} C_\alpha \cap D_\alpha = A \cap B \in \mathcal{F}_\alpha \dots (b), \\ (a), (b) &\Rightarrow (\forall \alpha \in I_0)(\exists (X, \mathcal{E}) \text{ t.s.}) (\mathcal{F}_\alpha \text{ base for } \mathcal{E}). \end{aligned}$$

2. $(\mathcal{F} \subset I^X)(|\mathcal{F}| < \aleph_0) \Rightarrow [(X, \mathcal{F}) \text{ f.t.s.} \Leftrightarrow (\forall \alpha \in I_0)(X, \mathcal{F}_\alpha) \text{ t.s.}]$

Proof : i) \Rightarrow :

$$\begin{aligned} (X, \mathcal{F}), \text{ f.t.s.} &\stackrel{2.3}{\Rightarrow} (0 \in \mathcal{F})(1 \in \mathcal{F}) \\ &\stackrel{2.6-2.7}{\Rightarrow} (\forall \alpha \in I_0)(0_\alpha = \emptyset \in \mathcal{F}_\alpha)(1_\alpha = X \in \mathcal{F}_\alpha) \dots (a), \\ A, B \in \mathcal{F}_\alpha &\stackrel{4.1}{\Rightarrow} A \cap B \in \mathcal{F}_\alpha \dots (b), \\ A \subset \mathcal{F}_\alpha &\stackrel{2.7.}{\Rightarrow} (\exists B \subset \mathcal{F})(A = B_\alpha) \\ &\left. \begin{array}{l} \mathcal{F} \text{ f.t.} \\ \left. \begin{array}{l} \stackrel{2.7}{\Rightarrow} (\forall B)_\alpha \in \mathcal{F}_\alpha \stackrel{\text{hip.}-3.5.}{\Rightarrow} \bigcup B_\alpha = \bigcup A \in \mathcal{F}_\alpha \dots (c), \end{array} \right\} \Rightarrow \forall B \in \mathcal{F} \end{array} \right\} \end{aligned}$$

$$\begin{aligned}
 & \text{(a),(b),(c)} \Rightarrow (\alpha \in I_0 \Rightarrow (X, \mathcal{F}_\alpha) \text{ t.s.}), \\
 & \text{ii)} \Rightarrow: (X, \mathcal{F}) \text{ not f.t.s.} \\
 & \quad \xrightarrow{2.4.} \Rightarrow (0 \notin \mathcal{F}) \vee (1 \notin \mathcal{F}) \vee (\exists A, B \in \mathcal{F})(A \wedge B \notin \mathcal{F}) \vee (\exists \mathcal{A} \subset \mathcal{F})(\bigvee \mathcal{A} \notin \mathcal{F}) \dots \text{(a)}, \\
 & \mathbf{0} \notin F \Rightarrow (\forall A \in \mathcal{F})(\exists a \in X)(0 < A(a) \leq 1) \\
 & \quad \left. \begin{array}{l} \alpha = A(a)/2 \\ (\exists \alpha \in I_0)(\forall A \in \mathcal{F})(\exists a \in X)(A(a) > \alpha) \Rightarrow \\ (\exists \alpha \in I_0)(\forall A \in \mathcal{F})(\exists a \in A_\alpha) \Rightarrow (\exists \alpha \in I_0)(\phi \notin \mathcal{F}_\alpha) \dots \text{(b)}, \end{array} \right\} \Rightarrow \\
 & \mathbf{1} \notin F \Rightarrow (\forall A \in \mathcal{F})(\exists x \in X)(0 \leq A(x) < 1) \\
 & \quad \Rightarrow (\forall A \in \mathcal{F})(A_1 \neq X) \Rightarrow (\exists \alpha \in I_0)(X \notin \mathcal{F}_\alpha) \dots \text{(c)}, \\
 & (\exists A, B \in \mathcal{F})(A \wedge B \notin \mathcal{F}) \xrightarrow{3.1} \\
 & \quad [\exists \alpha \in I_0 \setminus \{ \alpha \mid (\exists C \in \mathcal{F})((A \wedge B)_x = C_x)(A \wedge B \neq C) \}] \\
 & \quad \quad \quad (A_\alpha, B_{\alpha \in \mathcal{F}_\alpha})(A \wedge B)_\alpha \notin \mathcal{F}_\alpha \\
 & \quad \quad \quad \xrightarrow{3.2} \Rightarrow (\exists \alpha \in I_0)(A_\alpha, B_{\alpha \in \mathcal{F}_\alpha})(A_\alpha \cap B_\alpha \notin \mathcal{F}_\alpha) \dots \text{(d)}, \\
 & (\exists \mathcal{A} \subset \mathcal{F})(\bigvee \mathcal{A} \notin \mathcal{F}) \xrightarrow{3.1} \\
 & \quad K = I \setminus \{ x \mid (\exists C \in \mathcal{F})((\bigvee \mathcal{A})_x = C_x)(\bigvee \mathcal{A} \neq C) \} \neq \emptyset \\
 & \quad \quad \quad \left. \begin{array}{l} \alpha \in K \end{array} \right\} \Rightarrow \\
 & \quad \quad \quad (\exists \alpha \in I_0)(\mathcal{A}_\alpha \subset \mathcal{F}_\alpha)(\bigvee \mathcal{A})_\alpha \notin \mathcal{F}_\alpha \xrightarrow{\text{Hip-3.5}} \\
 & \quad \quad \quad (\exists \mathcal{B} = \mathcal{A}_\alpha \subset \mathcal{F}_\alpha)(\bigcup \mathcal{B} = \bigcup \mathcal{A}_\alpha \notin \mathcal{F}_\alpha) \dots \text{(e)}, \\
 & \text{(a),(b),(c),(d), (e)} \Rightarrow (\exists \alpha \in I_0) [(X, \mathcal{F}_\alpha) \text{ not t.s. }].
 \end{aligned}$$

Example:

$$\begin{aligned}
 & X = (0, 1), \quad A = \{ (x, -x/6 + 2/3) \mid x \in X \}, \quad B = \{ (x, x/2 + 1/4) \mid x \in X \}, \\
 & \mathcal{F} = \{ \mathbf{0}, \mathbf{1}, A, B, A \vee B, A \wedge B \} \Rightarrow |\mathcal{F}| = 6 < \aleph_0, \\
 & (A \vee B)(x) = \begin{cases} x/2 + 1/4, & 0 < x \leq 5/8 \\ -x/6 + 2/3, & 5/8 < x < 1 \end{cases}, \quad (A \wedge B)(x) = \begin{cases} -x/6 + 2/3, & 0 < x \leq 5/8 \\ x/2 + 1/4, & 5/8 < x < 1 \end{cases} \\
 & A_\alpha = \begin{cases} X, & 0 < \alpha \leq 1/2 \\ (0, 4 - 6\alpha], & 1/2 < \alpha < 2/3 \\ \phi, & 2/3 \leq \alpha \leq 1 \end{cases}, \quad B_\alpha = \begin{cases} X, & 0 < \alpha \leq 1/4 \\ [2\alpha - 1/2, 1), & 1/4 < \alpha < 3/4 \\ \phi, & 3/4 \leq \alpha \leq 1 \end{cases} \\
 & (A \vee B)_\alpha = A_\alpha \vee B_\alpha = \begin{cases} X, & 0 < \alpha \leq 9/16 \\ (0, 4 - 6\alpha] \cup [2\alpha - 1/2, 1), & 9/16 < \alpha < 2/3 \\ [2\alpha - 1/2, 1), & 2/3 \leq \alpha < 3/4 \\ \phi, & 3/4 \leq \alpha \leq 1 \end{cases} \\
 & (A \wedge B)_\alpha = A_\alpha \wedge B_\alpha = \begin{cases} X, & 0 < \alpha \leq 1/4 \\ [2\alpha - 1/2, 1), & 1/4 < \alpha < 1/2 \\ [2\alpha - 1/2, 4 - 6\alpha], & 1/2 \leq \alpha \leq 9/16 \\ \phi, & 9/16 < \alpha \leq 1 \end{cases}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{F}_\alpha &= \{ \mathbf{0}_\alpha, \mathbf{1}_\alpha, A_\alpha, B_\alpha, (A \vee B)_\alpha, (A \wedge B)_\alpha \} = \{ \phi, X, A_\alpha, B_\alpha, A_\alpha \vee B_\alpha, A_\alpha \wedge B_\alpha \} \\
 &= \begin{cases} \{ \phi, X \}, & 0 < \alpha \leq 1/4 \vee 3/4 < \alpha \leq 1 \\ \{ \phi, X, [2\alpha-1/2, 1) \}, & 1/4 < \alpha \leq 1/2 \vee 2/3 < \alpha < 3/4 \\ \{ \phi, X, [2\alpha-1/2, 1), (0, 4-6\alpha], [2\alpha-1/2, 4-6\alpha] \}, & 1/2 < \alpha \leq 9/16 \\ \{ \phi, X, [2\alpha-1/2, 1), (0, 4-6\alpha], (0, 4-6\alpha] \cup [2\alpha-1/2, 1) \}, & 9/16 < \alpha < 2/3 \end{cases} \\
 & \quad (\forall \alpha \in I_0) (X, \mathcal{F}_\alpha) \text{ t.s. } .
 \end{aligned}$$

3. $\{ \cup \mathcal{A} \mid \mathcal{A} \subset \mathcal{F} \} \subset \mathcal{F} \subset \mathcal{I}^X \Rightarrow [(X, \mathcal{F}) \text{ f.t.s.} \Leftrightarrow (\forall \alpha \in I_0) (X, \mathcal{F}_\alpha) \text{ t.s.}]$

Proof : from 3.6., 4.1. and 4.2.

4. $(\exists (X, \mathcal{F}) \text{ f.t.s.}) (\exists \alpha \in I_0) (X, \mathcal{F}_\alpha) \text{ not t.s. }$

Proof : (Example)

$$X = \mathbb{R}, \quad A^a(x) = \begin{cases} 0, & x < a \\ x-a, & a \leq x \leq a+1 \\ 1, & a+1 < x \end{cases} \quad F = \{ 0, 1 \} \cup \{ A^a \mid a \geq 0 \} \Rightarrow$$

$\exists (X, \mathcal{F}) \text{ f.t.s. }$, but for $\alpha = 1/2$, $\mathcal{F}_\alpha = \{ \phi, \mathbb{R} \} \cup \{ [a+1/2, \infty) \mid a \geq 0 \}$ and $\mathcal{A} = \{ [a+1/2, \infty) \mid a > 0 \} \subset F_\alpha$, $\cup \mathcal{A} = (1/2, \infty) \notin \mathcal{F}_\alpha$ and $(\mathbb{R}, \mathcal{F}_\alpha)$ not t.s.

5. $(\exists X) (\exists \mathcal{A} \subset \mathcal{I}^X) (\forall \alpha \in I_0) (X, \mathcal{A}_\alpha) \text{ t.s.} (X, \mathcal{A}) \text{ not f.t.s. })$

Proof : (Example)

$$X = \mathbb{R}, \quad A^b = \{ (x, b) \mid x \in \mathbb{R} \}, \quad \mathcal{A} = \{ 0, 1 \} \cup \{ A^b \mid 0 < b < 1/2 \}, \quad \alpha \in I_0 \Rightarrow \\
 \mathcal{A}_\alpha = \{ 0, 1 \}, \quad (X, \mathcal{A}_\alpha) \text{ t.s.}, \text{ but } (X, \mathcal{A}) \text{ not f.t.s. } .$$

6. $(X, \mathcal{F}) \text{ f.t.s.}) (\alpha \in I_0) (X, \mathcal{F}_\alpha) \text{ t.s.}) (A \subset X) \Rightarrow (A^\circ)_\alpha \subset (A_\alpha)^\circ$

Proof :

$$\left. \begin{array}{l} A^\circ \leq A \quad \xrightarrow{3.1.} \Rightarrow (A^\circ)_\alpha \subset A_\alpha \\ A^\circ \in F \quad \xrightarrow{2.7.} \Rightarrow (A^\circ)_\alpha \in \mathcal{F}_\alpha \end{array} \right\} \xrightarrow{\text{hip.}} \Rightarrow (A^\circ)_\alpha \subset (A_\alpha)^\circ$$

7. $(X, \mathcal{F}) \text{ f.t.s.}) (\alpha \in I_0) (X, \mathcal{F}_\alpha) \text{ t.s.}) (A \in \mathcal{F}) \Rightarrow (A^\circ)_\alpha = (A_\alpha)^\circ$

Proof.:

$$\left. \begin{array}{l} A \in \mathcal{F} \Rightarrow A^\circ = A \quad \xrightarrow{3-1.} \Rightarrow (A^\circ)_\alpha = A_\alpha \\ A \in \mathcal{F} \Rightarrow A \in \mathcal{F}_\alpha \quad \xrightarrow{2.7.} \Rightarrow (A_\alpha)^\circ = A_\alpha \end{array} \right\} \xrightarrow{\text{hip.}} \Rightarrow (A^\circ)_\alpha = (A_\alpha)^\circ$$

8. $(X, \mathcal{F}) \text{ f.t.s.}) (Y, \mathcal{E}) \text{ f.t.s.})$

$$(\alpha \in I_0) (X, \mathcal{F}_\alpha) \text{ t.s.}) (Y, \mathcal{E}_\alpha) \text{ t.s.}) (f : (X, \mathcal{F}) \rightarrow (Y, \mathcal{E}) \text{ f.c.}) \\
 \Rightarrow f : (X, \mathcal{F}_\alpha) \rightarrow (Y, \mathcal{E}_\alpha) \text{ c.}$$

Proof.:

$$\begin{array}{ll} B \in \mathcal{E}_\alpha \xrightarrow{2.7.} (\exists A \in \mathcal{E}) (B = A_\alpha) & \xrightarrow{\text{Hip-2.10-}} f^1[A] \in \mathcal{F} \\ \xrightarrow{2.7.} (f^1[A])_\alpha \in \mathcal{F}_\alpha & \xrightarrow{3.9.} f^1[A_\alpha] = f^1[B] \in \mathcal{F}_\alpha \end{array} \\
 / \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad f : (X, \mathcal{F}_\alpha) \rightarrow (Y, \mathcal{E}_\alpha) \text{ c.}$$

$$9. \quad f : (X, \mathcal{F}) \rightarrow (Y, \mathcal{E}) \text{ f.c.} \Leftrightarrow (\forall A \in I^Y) (f^{-1}[A^0] \leq (f^{-1}[A])^0)$$

Proof.: i. \Rightarrow :

$$\left. \begin{array}{l} A \in I^Y \Rightarrow A^0 \in \mathcal{E} \xrightarrow{\text{Hip.}} f^{-1}[A^0] \in \mathcal{F} \\ A^0 \leq A \Rightarrow f^{-1}[A^0] \leq f^{-1}[A] \end{array} \right\} \xrightarrow{2.4} f^{-1}[A^0] \leq (f^{-1}[A])^0$$

ii. \Leftarrow :

$$\begin{aligned} A \in \mathcal{E} &\Rightarrow A^0 = A \xrightarrow{2.4} f^{-1}[A^0] = f^{-1}[A] \\ &\xrightarrow{\text{hip.}} f^{-1}[A] \leq (f^{-1}[A])^0 \xrightarrow{2.4} f^{-1}[A] = (f^{-1}[A])^0 \\ &\xrightarrow{2.4} f^{-1}[A] \in \mathcal{F} \end{aligned}$$

/ f f.c.

$$10.. \quad ((X, \mathcal{F}) T_0) (\beta \in I_0) ((X, \mathcal{F}_\beta) \text{ t.s}) \Rightarrow (X, \mathcal{F}_\beta) T_0$$

Proof.:

$$\begin{aligned} (X, \mathcal{F}) T_0 &\xrightarrow{2.10} \\ (\forall x, y \in X) (\forall \alpha \in I_0) (\exists A \in \mathcal{F}) [(P_x^\alpha \in A) (P_y^\alpha \notin A) \vee (P_x^\alpha \notin A) (P_y^\alpha \in B)] \\ &\xrightarrow{2.7} (\forall x, y \in X) (A_\beta \in \mathcal{F}_\beta) [(x \in A_\beta) (y \notin A_\beta) \vee (x \notin A_\beta) (y \in B_\beta)] \end{aligned}$$

$$\begin{aligned} &\Rightarrow (X, \mathcal{F}) T_0 \\ \text{Example: } X = \{ a, b \}, \mathcal{F} = \{ 0, 1, P_a^{1/3} \}, A = P_a^{1/3} \\ A(x) = \begin{cases} 0, & x \neq a \\ 1/3, & x = a \end{cases}, \text{ for } \alpha = 1/4, A_\alpha = \{a\}, \mathcal{F}_\alpha = \{ \phi, X, \{a\} \} T_0 \end{aligned}$$

Concluding Remarks

In this study, various basic theorems on fuzzy sets and their α -cross-sections, and topological fuzzy-crisp relations resulting from these theorems have been introduced. Similarly, many fuzzy-topological concepts can be related with their equivalent crisp-topological concepts.

In the proofs presented here, pure symbolic and formal language of mathematics, which is a universal mean of communication, has been deliberately used instead of a national language.

References:

- [1] Hu Cheng- Ming, Fuzzy Topological Spaces, Journal of Mathematical Analysis and Applications, 110, 141-178, (1985)
- [2] Chang, C.L., Fuzzy Topological Spaces, Journal of Mathematical Analysis and Applications, 24, 182-190, (1968)
- [3] Husain, T., Almost Continuous Mappings. Prace Math., 10, 1-7(1966)
- [4] Güney, Z., Fuzzy Kümelerin α - kesitlerine dair bazı sonuçlar ve bir düzeltme, Ulusal Matematik Sempozyumu, Çukurova Üniv. Adana, 1995